



Lipschitz stability of canonical Jordan bases of H -selfadjoint matrices under structure-preserving perturbations

T. Bella *, V. Olshevsky, U. Prasad

Department of Mathematics, University of Connecticut, Storrs CT 06269-3009, USA

Received 21 May 2007; accepted 19 November 2007

Available online 5 February 2008

Submitted by E. Tyrtyshnikov

Abstract

In this paper we study Jordan-structure-preserving perturbations of matrices selfadjoint in the *indefinite inner product*. The main result of the paper is Lipschitz stability of the corresponding *affiliation* matrices. The result can be reformulated as Lipschitz stability, under small perturbations, of *canonical* Jordan bases (i.e., eigenvectors and generalized eigenvectors enjoying a certain *flipped* orthonormality relation) of matrices selfadjoint in the indefinite inner product. The proof relies upon the analysis of small perturbations of invariant subspaces, where the size of a permutation of an invariant subspace is measured using the concepts of a *gap* and of a *semigap*.

© 2007 Published by Elsevier Inc.

Keywords: Indefinite inner product; Perturbations; Structure-preserving perturbations; Canonical Jordan bases; Invariant subspaces; Gaps

1. Introduction. Part I. Preliminaries

1.1. Motivation and main result

Perturbation problems for matrices have been studied by many authors in different contexts, see, e.g., the monographs [5,17,1,9] among others, as well as the references therein. To motivate

* Corresponding author.

E-mail addresses: bella@math.uconn.edu (T. Bella), olshevsky@math.uconn.edu (V. Olshevsky), upendra@math.uconn.edu (U. Prasad).

Table 1

A selection of motivating results on the perturbation of Jordan structure

	General perturbations (possibly changing Jordan structure)	Perturbations preserving Jordan structure
General matrices	(i) [3,12,2,11]	(ii) [7,13]
Matrices selfadjoint in indefinite inner product	(iii) [14]	(iv) [4,16], this paper

the problem considered in this paper we briefly recall several relevant results captured by the four cells (i)–(iv) of Table 1.

Again, there is a vast literature on the subject, and the selection in the above table is clearly far from being comprehensive. It includes several references that directly motivate the problem considered.

- (i) *General matrices. General perturbations.* It is well known that even small perturbations of a given matrix A_0 can destroy its Jordan structure. For instance, for a nearby matrix A , not only the eigenvalues of A but also the sizes of its corresponding Jordan blocks can be different from those of A_0 . For a fixed A_0 , the full description of all possible Jordan structures of nearby matrices A was conjectured by Gohberg and Kaashoek in [3]. It was proven independently in [12,2]. Two more proofs of the Gohberg–Kaashoek conjecture can be found in [11]. We do not discuss their general results in detail since in this paper we limit our focus to the special cases considered next.
- (ii) *General matrices. Perturbations preserving Jordan structure.* In [7] (see also [5]) the authors considered special perturbations A that *preserve the Jordan structure* of A_0 . We start with the following simplified version of their result.

Proposition 1.1 (Lipschitz stability of similarity matrices). *Let $A_0 \in \mathbb{C}^{n \times n}$ be a fixed matrix. There is a constant $K > 0$ (depending on A_0 only) such that for any A that is similar to A_0 there exists a similarity matrix S , i.e., $S^{-1}A_0S = A$, such that*

$$\|I - S\| \leq K \|A - A_0\|. \quad (1.1)$$

In words, if a small perturbation A of A_0 is similar to A_0 , then a (highly nonunique) similarity matrix S can be chosen to be a small perturbation of the identity matrix, and a Lipschitz-type bound (1.1) holds.

In fact, Gohberg and Rodman considered more general perturbations A that are *not similar* to A_0 but *have the same Jordan structure*. Since the latter concept plays a key role in what follows we give two relevant definitions next.

Definition 1.2

- (*Same Jordan structure*). Denote by $\sigma(A_0)$ and $\sigma(A)$ the sets of all eigenvalues of A_0 and A , respectively. Matrices A_0 and A are said to have *the same Jordan structure* if there is a bijective function $f : \sigma(A_0) \rightarrow \sigma(A)$ such that if $\mu = f(\lambda)$, then λ and μ have the same Jordan block sizes.

- (Same Jordan bases). Matrices A_0 and A that have the same Jordan structure are said to additionally have *the same Jordan bases* if the following statement is true. If $\mu = f(\lambda)$, then every Jordan chain of A_0 corresponding to λ is also a Jordan chain of A corresponding to μ (and automatically vice versa).

Remark 1.3 (Same Jordan bases). Two matrices $A_0, A \in \mathbb{C}^{n \times n}$ have the same Jordan bases if the following statement holds. If, for an invertible T , the matrix $T^{-1}A_0T$ is in a canonical Jordan form, then $T^{-1}AT$ is also in a canonical Jordan form.

In order to generalize Proposition 1.1 to perturbations A having the same Jordan structure as A_0 we need to extend the concept of a similarity matrix S . The following obvious result is an enabling tool for doing this.

Lemma 1.4 (Affiliation matrix). Two matrices $A_0, A \in \mathbb{C}^{n \times n}$ have the same Jordan structure if and only if there is an invertible matrix S such that A_0 and $S^{-1}AS$ have the same Jordan bases.

We suggest to refer to the matrix S in Lemma 1.4 as an *affiliation matrix* since it generalizes the similarity matrix to the situation when A_0 and A might not be similar but have the same Jordan structure. Observe that an affiliation matrix is highly nonunique (just as its special case, a similarity matrix). We are now ready to present the following generalization of Proposition 1.1 that is implicit in [7,5].

Proposition 1.5 (Lipschitz stability of affiliation matrices). Let $A_0 \in \mathbb{C}^{n \times n}$ be a fixed matrix. There is a constant $K > 0$ (depending on A_0 only) such that for any A having the same Jordan structure as A_0 there exists an affiliation matrix S (i.e., matrices A_0 and $S^{-1}AS$ have the same Jordan bases), such that

$$\|I - S\| \leq K \|A - A_0\|. \tag{1.2}$$

- (iii) *H-selfadjoint matrices. General perturbations.* Matrices and their perturbations considered in the items (i) and (ii) above were general. It is of interest to study situations when both matrices A_0 and A have some special structure. Hermitian structure is of little interest in the context of perturbations of Jordan structure since Hermitian (or selfadjoint) matrices are diagonalizable and they cannot have Jordan blocks of size greater than one. Matrices that are selfadjoint with respect to an indefinite inner product appear in a number of applications [6], and they can have nontrivial Jordan blocks, so perturbation problems for their Jordan structure are of interest.

We refer to [6] for a comprehensive introduction to the subject, and only recall here that for a Hermitian, invertible (not necessarily positive definite) matrix H , one defines the *indefinite inner product* by

$$[x, y]_H = (Hx, y) = y^* Hx, \quad \text{where } (x, y) = y^* x \text{ is the standard Euclidean inner product.} \tag{1.3}$$

Further, a matrix is *H-selfadjoint* (or selfadjoint with respect to H) if

$$[Ax, y]_H = [x, Ay]_H \quad (\text{for all } x, y \in \mathbb{C}^n) \quad \text{or, equivalently, } HA = A^* H. \tag{1.4}$$

Note that setting $H = I$ in (1.3) and (1.4), one obtains the standard Euclidean inner product (\cdot, \cdot) , and standard selfadjoint (or Hermitian) matrices A .

The monograph [4] contains a number of results on the perturbation of eigenvalues of H -selfadjoint matrices. The variation of the Jordan structure of H -selfadjoint matrices under small perturbations was studied in [14] where one can find certain restrictions additional to those of [3,12,2,11] mentioned in the item (i) above. The techniques used in [14] allow us to obtain an analog of Proposition 1.5 for H -selfadjoint matrices, which is described next.

- (iv) *Main result. H -selfadjoint matrices. Perturbations preserving Jordan structure.* Let H_0 be a fixed invertible Hermitian matrix, and let A_0 be a fixed H_0 -selfadjoint matrix. We consider their perturbations A and H where A is H -selfadjoint (in particular, H is invertible and Hermitian). This case was considered in [4,16] where a number of results were obtained (we use some of them below). However, it seems the question of finding an analog of Proposition 1.5 has not been addressed in the literature yet. In order to obtain such an analog below one needs to carry over the concept of an affiliation matrix S (appearing in the Lipschitz-type bound (1.2)) to perturbations of matrices selfadjoint with respect to an indefinite inner product. The problem is that for an H -selfadjoint matrix A , a similar matrix $S^{-1}AS$ is not necessarily H -selfadjoint. This suggests that (in order to preserve the property of A of being selfadjoint with respect to indefinite inner product) the matrix H should also be modified appropriately. Here is the recipe.

Definition 1.6 (*Affiliation relation*). Let A_0 be H_0 -selfadjoint and A be H -selfadjoint.

- We will use the notations

$$(A, H) \xrightarrow{S} (A_0, H_0) \quad \text{to mean that} \quad S^{-1}AS = A_0 \quad \text{and} \quad S^*HS = H_0. \tag{1.5}$$

- The relation “ $(\cdot, \cdot) \xrightarrow{S} (\cdot, \cdot)$ ” will be called the *affiliation relation* of matrices (A, H) , where A is H -selfadjoint.

Two remarks are due.

- A simple calculation shows that this notation makes sense; i.e., if A is H -selfadjoint and $(A, H) \xrightarrow{S} (B, G)$, then B is G -selfadjoint.
- It is easy to see that affiliation is an equivalence relation.

In the above definition the matrices A_0 and A are similar, so the corresponding S was indeed a similarity matrix. In the following definition we consider the case when A_0 and A only have the same Jordan structure, and specify the concept of the affiliation matrix for the framework of an indefinite inner product.

Definition 1.7 (*Weak affiliation matrix*). Let A_0 be H_0 -selfadjoint and A be H -selfadjoint.

- A matrix S is called a (*weak*) *affiliation matrix* of the quadruple (A_0, H_0, A, H) if

$$(A, H) \xrightarrow{S} (A_1, H_0),$$

where matrices A_0 and A_1 have the same Jordan bases.

- In this case the pairs (A_0, H_0) and (A, H) are called (*weakly*) *affiliated*.

With this background we can introduce the main result proved in the paper.

Theorem 1.8 (Main result. Lipschitz stability of affiliation matrices). *Let $A_0 \in \mathbb{C}^{n \times n}$ be a fixed H_0 -selfadjoint matrix. There exist constants $K, \delta > 0$ (depending on A_0 and H_0 only) such that the following assertion holds. For any H -selfadjoint matrix A such that A has the same Jordan structure as A_0 and*

$$\|A - A_0\| + \|H - H_0\| < \delta, \tag{1.6}$$

the pairs (A_0, H_0) and (A, H) are affiliated, and there exists an affiliation matrix S such that

$$\|I - S\| \leq K(\|A - A_0\| + \|H - H_0\|). \tag{1.7}$$

In words, if (a) a small perturbation A of A_0 has the same Jordan structure as A_0 ; (b) H is a small perturbation of H_0 ; (c) A_0 is H_0 -selfadjoint and A is H -selfadjoint, then (i) an affiliation matrix S exists, and (ii) it can be chosen to be a small perturbation of the identity matrix, and a Lipschitz-type bound (1.7) holds.

Remark 1.9. There is a certain (deliberate) controversy here: while Definition 1.7 introduced a *weak* affiliation matrix, the Theorem 1.8 asserts the existence of a (*strong*) affiliation matrix to be formally introduced only in Definition 8.5 of Section 8. The controversy is only virtual since as we will see in Section 8, the weak affiliation matrix S constructed in the course of proof of Theorem 1.8 will enjoy several additional nice properties. Hence the (strong) affiliation matrix will be defined as a weak affiliation matrix having those additional properties. For this reason, in Sections 2–7 the term “affiliation” will be tentatively understood in the *weak* sense. In Section 8 it will be justified that all the results of the paper including Theorem 1.8 remain valid if the term “affiliation” is understood in the *strong* sense. Therefore we will just use the nomenclature “affiliation” without specifying whether it is weak or strong.

We preferred to formulate Theorem 1.8 before Definition 8.5 since the latter requires introducing a number of (unnecessary at the moment) technical details that will be dealt with in Sections 2 and 8.

Comparing Proposition 1.5 and Theorem 1.8, we see that the latter uses the assumption (1.6) not appearing in the former. We conclude this subsection with a simple example indicating that the condition (1.6) is essential, and it cannot simply be omitted.

Example 1.10 (*Affiliation matrix may not exist for large perturbations*). Let us consider 1×1 case:

$$A_0 = [1], \quad H_0 = [1], \quad A = [1], \quad H = [-1].$$

In this case the desired affiliation matrix S does not exist, since it must satisfy $S^* \cdot 1 \cdot S = -1$. Clearly, 1×1 matrices A_0 and A always have the same Jordan structure. However, H_0 and H are not close enough to ensure that (1.6) yields (1.7). In Section 2.1 we will recall another explanation [6] of the fact that S does not exist here, it will be based on the concept of the so-called *sign characteristic* whose definition is recalled in Section 2.

1.2. Structure of the paper

The next section continues the introduction with three interpretations of Theorem 1.8, one of which is the second main result of the paper, Theorem 2.6. The section concludes with a

graphical representation of the flow of the proofs of the paper. Section 3 presents a theorem showing it is sufficient for the proof of Theorem 1.8 to obtain the result for all pairs (A_0, H_0) in the canonical form. In Section 4, Theorem 1.8 is proved in the case where A_0 consists of a single Jordan block corresponding to a real eigenvalue, or a pair of Jordan blocks corresponding to a single nonreal eigenvalue. Following this, Section 5 presents a decoupling result that allows the process of Section 4 to apply inductively. The proof of this result requires some auxiliary results on semigaps and gaps between subspaces which are given in Section 6. These results are then used to prove the results of Section 5 in Section 7. In Section 8, the second main result of the paper, Theorem 2.6 is proved, and the details of the distinction between weak and strong affiliation introduced in Definition 1.7 and Remark 1.9 are explained in detail. Finally, in Section 9, the results of Theorems 1.8 and 2.6 are extended to the case of perturbations that *partially* preserve Jordan structure; that is, the sizes of Jordan blocks corresponding to some subset of the eigenvalues are unchanged.

2. Introduction. Part II. Three interpretations of Theorem 1.8

The second part of the introduction is somewhat more technical. In Sections 2.2 and 2.3 below we provide three useful interpretations of our main result. They will use two key concepts defined next.

2.1. Key definitions. Sign characteristic and canonical Jordan bases

We begin by quoting a fundamental theorem of [18,19] (see also [10,6]) which plays a central role in all arguments below. As usual, $J(\lambda)$ denotes a single Jordan block of the form

$$J(\lambda) = \begin{bmatrix} \lambda & 1 & 0 & \cdots & 0 \\ 0 & \lambda & 1 & \ddots & \vdots \\ \vdots & \ddots & \lambda & \ddots & 0 \\ \vdots & & & \ddots & 1 \\ 0 & \cdots & \cdots & 0 & \lambda \end{bmatrix}, \quad \tilde{J}(\lambda) = \begin{cases} J(\lambda) & \lambda \in \mathbb{R} \\ \begin{bmatrix} J(\lambda) & 0 \\ 0 & J(\bar{\lambda}) \end{bmatrix} & \lambda \notin \mathbb{R} \end{cases},$$

$$\tilde{I} = \begin{bmatrix} 0 & \cdots & \cdots & 0 & 1 \\ \vdots & & 0 & 1 & 0 \\ \vdots & \ddots & \ddots & 0 & \vdots \\ 0 & \ddots & \ddots & & \vdots \\ 1 & 0 & \cdots & \cdots & 0 \end{bmatrix}. \tag{2.1}$$

The matrix \tilde{I} was called the sip (standard involutory permutation) matrix in [5].

Theorem 2.1 ([18,19]. Canonical form of matrices selfadjoint with respect to indefinite inner product).

Let $A \in \mathbb{C}^{n \times n}$ be a fixed H -selfadjoint matrix for some invertible, selfadjoint matrix H . Then there exists an invertible matrix T such that

$$(A, H) \xrightarrow{T} (J, P), \quad (2.2)$$

where

$$J = J(\lambda_1) \oplus \cdots \oplus J(\lambda_\alpha) \oplus \tilde{J}(\lambda_{\alpha+1}) \oplus \cdots \oplus \tilde{J}(\lambda_\beta) \quad (2.3)$$

is a Jordan normal form of A for real eigenvalues $\lambda_1, \dots, \lambda_\alpha$ and nonreal eigenvalues $\lambda_{\alpha+1}, \dots, \lambda_\beta$ from the upper half-plane, and

$$P = P_1 \oplus \cdots \oplus P_\alpha \oplus P_{\alpha+1} \oplus \cdots \oplus P_\beta, \quad (2.4)$$

where P_k is a signed sip matrix $\epsilon_k \tilde{I}$ of the same size as $J(\lambda_k)$ (for $k = 1, \dots, \alpha$), and a sip matrix \tilde{I} of the same size as $\tilde{J}(\lambda_k)$ (for $k = \alpha + 1, \dots, \beta$), and $\epsilon_k = \pm 1$ for $k = 1, \dots, \alpha$. The set

$$\epsilon = \{\epsilon_1, \dots, \epsilon_\alpha\} \quad (2.5)$$

is determined by the pair (A, H) uniquely, up to a permutation of the signs ϵ_k corresponding to Jordan blocks of the same size and of the same eigenvalue.

The proof of this theorem can be found, for instance, in [6, Theorem 5.1.1].

Remark 2.2 (*Symmetry of eigenvalues*). In particular, nonreal eigenvalues of an H -selfadjoint matrix A come in complex conjugate pairs. Furthermore, for each nonreal conjugate pair of eigenvalues the sizes of their Jordan blocks are identical.

Definition 2.3 (*Canonical form and sign characteristic* [4]). The pair (J, P) in (2.3) and (2.4) is called a canonical form of (A, H) . The set of signs in (2.5) is called the *sign characteristic* of the pair (A, H) .

Recall that affiliation is an equivalence relation, and hence pairs that have different canonical forms (up to an appropriate rearrangement of Jordan blocks of A and corresponding blocks of H) cannot be similar. This is exactly what happened in Example 1.10. Indeed, it is immediate to see that the pairs (A_0, H_0) and (A, H) are in the canonical form, from which we can see they have different sign characteristics, and therefore they can not be similar.

The first equation (2.2) implies $T^{-1}AT = J$, which means that the columns of the matrix T form a Jordan basis of A . However, not all such matrices T satisfy the second equation $T^*HT = P$, also implied by (2.2), with P of (2.4). We coin a special name for the columns of those matrices T that satisfy both equations implied by (2.2).

Definition 2.4 (*Canonical Jordan basis of an H -selfadjoint matrix*). Let A be an H -selfadjoint matrix, and let T be a similarity matrix that brings (A, H) to its canonical form (J, P) . The columns of T form a Jordan basis of A that will be called a canonical Jordan basis of (A, H) .

The following example makes the property of “flipped orthonormality” of the vectors of a canonical basis more transparent.

Example 2.5 (*Flipped orthonormality*). Let

$$J = \left[\begin{array}{ccc|cc} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right], \quad P = \left[\begin{array}{ccc|cc} 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & -1 & 0 \end{array} \right]$$

be a canonical pair. The canonical Jordan basis $\{\{e_1, e_2, e_3\}, \{e_4, e_5\}\}$ of (J, P) consists of two Jordan chains

$$0 \leftarrow e_1 \leftarrow e_2 \leftarrow e_3, \quad 0 \leftarrow e_4 \leftarrow e_5.$$

It is easy to see that the *canonical* structure of P yields that vectors belonging to different chains of this *canonical basis* are P -orthogonal, i.e.,

$$[e_j, e_k]_P = 0 \quad (j = 1, 2, 3; k = 4, 5).$$

Further, for the same reason the vectors within one chain have what might be called “flipped orthonormality,”:

$$[e_j, e_k]_P = \delta_{j,4-k} \quad (j, k = 1, 2, 3), \quad [e_j, e_k]_P = \delta_{j-3,3-(k-3)} \quad (j, k = 4, 5).$$

It is the above “flipped orthonormality property” that distinguishes canonical Jordan basis from the other ones.

With this background we are now ready to present a first interpretation of Theorem 1.8.

2.2. *First interpretation of the main result. Lipschitz stability of canonical Jordan bases of H-selfadjoint matrices under small perturbations preserving Jordan structure*

Let $\{\lambda_1, \dots, \lambda_\beta\}$ be a set of all eigenvalues of A_0 . Denote

$$m_k(A_0, \lambda_s) := \text{the size of the } k\text{th Jordan block of the matrix } A_0 \\ \text{corresponding to its eigenvalue } \lambda_s,$$

where the Jordan blocks are ordered such that their sizes are nonincreasing. Throughout the paper we assume that $\{m_k(A_0, \lambda_s)\}$ are ordered in nonascending order for each s . Let A have the same Jordan structure as A_0 , which means that the eigenvalues $\{\mu_1, \dots, \mu_\beta\}$ of A can be ordered such that

$$m_k(A_0, \lambda_s) = m_k(A, \mu_s).$$

With these notations Theorem 1.8 implies the following result on stability of eigenvectors and generalized eigenvectors.

Theorem 2.6 (*Lipschitz stability of canonical Jordan bases*). Let $A_0 \in \mathbb{C}^{n \times n}$ be a fixed H_0 -selfadjoint matrix. Let

$$\{ \{f_r^{(k,s)}\}_{r=0}^{m_k(A_0, \lambda_s)-1} \}_{s=1, k=1}^{s=\beta, k=\dim \text{Ker}(A_0 - \lambda_s I)} \tag{2.6}$$

be a fixed canonical Jordan basis of A_0 . There exist constants $K, \delta > 0$ (depending on A_0 and H_0 only) such that the following assertion holds. For any H -selfadjoint matrix A such that A has the same Jordan structure as A_0 and

$$\|A - A_0\| + \|H - H_0\| < \delta, \tag{2.7}$$

there exists a canonical Jordan basis

$$\{g_r^{(k,s)}\}_{r=0}^{m_k(A,\lambda_s)-1}\}_{s=1,k=1}^{s=\beta,k=\dim \text{Ker}(A-\lambda_s I)}$$

of A such that

$$\|g_r^{(k,s)} - f_r^{(k,s)}\| \leq K(\|A - A_0\| + \|H - H_0\|) \tag{2.8}$$

for all k, s, r within their ranges.

In words, let $\{f_r^{(k,s)}\}$ be a fixed canonical Jordan basis of a fixed H_0 -selfadjoint matrix A_0 , and let (A, H) be a small perturbation of (A_0, H_0) where A is H -selfadjoint. If A has the same Jordan structure as A_0 , then a (highly nonunique) canonical Jordan basis $\{g_r^{(k,s)}\}$ of (A, H) can be chosen to be a small perturbation of the given canonical Jordan basis of (A_0, H_0) , and the Lipschitz-type bound (2.8) holds.

The proof of the above result will be given later in Section 8.1.

2.3. Second interpretation of the main result. Affiliation matrix is (H_0, H) -unitary for small perturbations

Let us specify Theorem 1.8 to the case when $H_0 = H = I$. It is easy to see from the definition in (1.4) that in this case both matrices A_0 and A are Hermitian. Secondly, in this case (1.5) implies $S^*S = I$, i.e., S is a unitary matrix. To sum up, Theorem 1.8 specifies to the following result.

Theorem 2.7. *Let A_0 be a fixed Hermitian matrix. Then there exists a constant $K > 0$ such that for any Hermitian matrix A there exists a unitary affiliation matrix S (i.e., such that A_0 and $S^{-1}AS$ are simultaneously diagonalizable) such that*

$$\|I - S\| \leq K\|A - A_0\|. \tag{2.9}$$

In words, if A_0 and A are both Hermitian, then the affiliation matrix S can be chosen to be unitary and satisfying the bound (2.9). The latter result (Lipschitz stability of eigenvectors of Hermitian matrices) is known (e.g., it is an obvious consequence of [15]), but it leads to an interesting interpretation (cf. with [16]) of the main result of the paper, Theorem 1.8.

In this context, the meaning of Theorem 1.8 is that extending to the case of indefinite inner products, under the stated conditions the affiliation matrix S can be chosen to be (H_0, H) -unitary (i.e., $S^*HS = H_0$) and satisfying the Lipschitz-type bound (1.7).

2.4. Third interpretation of the main result. Lipschitz stability of congruency matrices

In Section 2.3 we considered a special case when the matrix H in (A, H) was the identity matrix, i.e., $H_0 = H = I$. Here we consider another special case and set $A_0 = A = I$. Clearly, I is H -selfadjoint for any invertible Hermitian H . Here is a specialization of our main result, Theorem 1.8, in this case.

Theorem 2.8 (Lipschitz stability of congruency matrices). *Let H_0 be a fixed invertible Hermitian matrix. There exist constants $K, \delta > 0$ (depending on H_0 only) such that the following assertion holds. For any Hermitian matrix H such that*

$$\|H - H_0\| < \delta, \tag{2.10}$$

there exists a congruency matrix S , i.e., $S^*HS = H_0$, such that

$$\|I - S\| \leq K \|H - H_0\|. \tag{2.11}$$


Note that in view of (2.10) and (2.11) both H and S are invertible, and $\|S\|$ and hence $\|S^{-1}\|$ are bounded. Hence

$$\|I - S^{-1}\| = \|S^{-1}(S - I)\| \leq \|S^{-1}\| \cdot \|I - S\| = K_1 \|H - H_0\|.$$

with some K_1 . This is a local version of the main result of [15] who proved the bound $\|I - S^{-1}\| \leq K_1 \|H - H_0\|$ without the restriction¹ (2.10), but requiring instead that H_0 and H remain congruent.

2.5. The flow of the results

The main results of the paper are Theorems 1.8 and 2.6. The following diagram presents the flow of the proofs.



The proof of Theorem 5.1 is the central point in establishing the main results, Theorems 1.8 and 2.6. As one can see in the diagram above, the proof of Theorem 5.1 is based on the results of Sections 4, 6, and 7.

¹ Note that examples indicate that without restriction (2.10) the bound (2.11) does not hold.

3. Reduction to the canonical form

We will find it useful throughout the paper to assume that the matrices A_0 and H_0 of Theorem 1.8 are in the canonical form. There is no loss of generality with this assumption, as the next theorem demonstrates.

Theorem 3.1 (Reduction to the canonical form). *Suppose the result of Theorem 1.8 is true for each pair (A_0, H_0) in the canonical form as defined in the Definition 2.3. Then the result of Theorem 1.8 is true for all pairs (B_0, G_0) , where B_0 is G -selfadjoint.*

Proof. Suppose the pair (B_0, G_0) is not in the canonical form. By Theorem 2.1, there exists a matrix T such that

$$(B_0, G_0) \xrightarrow{T} (A_0, H_0)$$

for some pair (A_0, H_0) in the canonical form. Then we have

$$\begin{array}{ccc} (A, H) & \xrightarrow{S} & (A_1, H_0) \\ T \uparrow & & T \uparrow \\ (B, G) & \xrightarrow{R=TSST^{-1}} & (B_1, G_0) \end{array} \tag{3.1}$$

The above diagram implies that the bound for a general (B_0, G_0) ,

$$\|I - R\| \leq K_{B_0, G_0} (\|B - B_0\| + \|G - G_0\|), \tag{3.2}$$

can be deduced from the bound for a *canonical* pair (A_0, H_0) ,

$$\|I - S\| \leq K_{A_0, H_0} (\|A - A_0\| + \|H - H_0\|).$$

Indeed, using the standard notation $\kappa(T) = \|T\| \cdot \|T^{-1}\|$ and the formulas captured by the diagram (3.1) we compute

$$\begin{aligned} \|I - R\| &= \|I - TST^{-1}\| \leq \kappa(T)\|I - S\| \leq \kappa(T)K_{A_0, H_0} (\|A - A_0\| + \|H - H_0\|) \\ &\leq \kappa(T)K_{A_0, H_0} (\|T^{-1}BT - T^{-1}B_0T\| + \|T^*GT - T^*G_0T\|) \\ &= \kappa(T)K_{A_0, H_0} (\kappa(T)\|B - B_0\| + \|T\|^2\|G - G_0\|) \\ &\leq K_{B_0, G_0} (\|B - B_0\| + \|G - G_0\|). \quad \square \end{aligned}$$

Hence it suffices to consider in what follows only the cases where (A_0, H_0) are in the canonical form.

4. Perturbations of a single real Jordan block or of a pair of complex conjugate Jordan blocks

In this section we present the first step of the proof of Theorem 1.8 for the special case where

$$A_0 = \tilde{J}(\lambda), \quad H_0 = \epsilon \tilde{I} \quad (\text{with } \epsilon = \pm 1), \tag{4.1}$$

where the sip matrix \tilde{I} and Jordan block $\tilde{J}(\lambda)$ were defined in (2.1). Recall that for a real λ the matrix $\tilde{J}(\lambda)$ is a single Jordan block, and for a nonreal λ we have $\epsilon = 1$ and the matrix $\tilde{J}(\lambda)$ is a direct sum of two Jordan blocks. In both cases it is easy to see that A_0 is H_0 -selfadjoint.

Theorem 4.1 (Lipschitz stability of affiliation matrices in the single block case). *Let $A_0 \in \mathbb{C}^{n \times n}$ be a fixed H_0 -selfadjoint matrix as given in (4.1). There exist constants $K, \delta > 0$ (depending on A_0 and H_0 only) such that the following assertion holds. For any H -selfadjoint matrix A such that A has the same Jordan structure as A_0 and*

$$\|A - A_0\| + \|H - H_0\| < \delta, \tag{4.2}$$

there exists an affiliation matrix S such that

$$\|I - S\| \leq K(\|A - A_0\| + \|H - H_0\|). \tag{4.3}$$

Theorem 4.1 will be proved in Section 4.2, and an extension of it to the case of a single complex eigenvalue will be discussed in Section 4.3. Before doing so, we illustrate three of the steps by which the proof will proceed with the following simple example.

4.1. Single-Jordan-block model example (for Theorem 4.1)

We begin this section with a simple example involving a matrix A_0 in Jordan form. For the pair

$$A_0 = \begin{bmatrix} 0 & 1 & 0 \\ & 0 & 1 \\ & & 0 \end{bmatrix} \quad \text{and} \quad H_0 = \begin{bmatrix} & & 1 \\ & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix},$$

consider the matrices

$$A = \begin{bmatrix} 0 & 1 & 2\delta \\ & 0 & 1 \\ & & 0 \end{bmatrix} \quad \text{and} \quad H = \begin{bmatrix} & & 1 \\ & 1 & -2\delta \\ 1 & -2\delta & \delta \end{bmatrix}$$

for some small positive δ . The pair (A, H) is then a small perturbation of the pair (A_0, H_0) , and it is straightforward to verify that A_0 is H_0 -selfadjoint and A is H -selfadjoint.

We wish to produce a matrix S satisfying²

- (i) $(A, H) \xrightarrow{S} (A_0, H_0)$;
- (ii) $\|I - S\| \leq K(\|A - A_0\| + \|H - H_0\|)$.

We will design this matrix S in three steps, the first will produce a matrix S_1 , the second will produce a matrix S_2 , and the third step will combine these as $S = S_1 S_2$, and check that it satisfies the desired bound.

4.1.1. First step. Mapping $A \rightarrow A_1$. Constructing S_1 .

Notice that the Jordan chain of A corresponding to $\lambda = 0$ is

$$0 \leftarrow e_1 \leftarrow e_2 + (2\delta)e_1 \leftarrow e_3,$$

where \leftarrow denotes application of the matrix A . This chain is a small perturbation of that of A_0 corresponding to $\lambda = 0$, which is simply

² Note that in this example, A and A_0 are similar (not the less restrictive condition of having the same Jordan structure), and hence $A_1 = J_3(0) = A_0$.

$$0 \leftarrow e_1 \leftarrow e_2 \leftarrow e_3.$$

The matrix that maps these basis vectors (those of A_0 into those of A) is given by

$$S_1 = \begin{bmatrix} 1 & 2\delta & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Then we have

$$S_1^{-1}AS_1 = A_0, \quad \text{and} \quad S_1^*HS_1 = \begin{bmatrix} & & 1 \\ 1 & 1 & 0 \\ 1 & 0 & \delta \end{bmatrix} =: H_1, \tag{4.4}$$

so that

$$(A, H) \xrightarrow{S_1} (A_0, H_1).$$

This illustrates the need for the second step below, as we need to generate an S such that $(A, H) \xrightarrow{S} (A_0, H_0)$ and $H_0 \neq H_1$.

4.1.2. *Second step. Zeroing sub-antidiagonal entries of H_1 . Constructing S_2 .*

The next step is to choose a matrix S_2 so that $(A_0, H_1) \xrightarrow{S_2} (A_0, H_0)$, that is, a matrix that repairs the problem below the anti-diagonal in H_1 of (4.4) to produce H_0 without modifying the fact that $S_1^{-1}AS_1$ already produced A_0 . The existence of such a matrix S_2 in general will be proven in the coming sections, but for now, notice that the matrix

$$S_2 = \begin{bmatrix} 1 & 0 & -\delta \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

satisfies the required conditions; that is

$$S_2^*H_1S_2 = \begin{bmatrix} & & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} = H_0 \quad \text{and} \quad S_2^{-1}A_0S_2 = S_2^{-1}S_2A_0 = A_0,$$

so that

$$(A_0, H_1) \xrightarrow{S_2} (A_0, H_0).$$

4.1.3. *Third step. Combining S_1 and S_2 .*

Set $S = S_1S_2$. We have demonstrated in the previous two steps that

$$(A, H) \xrightarrow{S_1} (A_0, H_1) \xrightarrow{S_2} (A_0, H_0) \quad \text{and hence} \quad (A, H) \xrightarrow{S} (A_0, H_0),$$

which implies condition (i). Furthermore, computing S explicitly yields

$$S = S_1S_2 = \begin{bmatrix} 1 & 2\delta & -\delta \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Hence all three differences $A - A_0$, $H - H_0$, and $I - S$ are of the same small order of δ , and so condition (ii) is satisfied.

4.2. Proof of Theorem 4.1 in the case of a real single Jordan block.

In this section we illustrate that the approach demonstrated in the above example can yield the proof of Theorem 4.1. The presentation follows that of the example in the previous section and is organized into three sections.

4.2.1. First step. Mapping $A \rightarrow A_1$. Constructing \tilde{S}_1 .

In this section we prove the following proposition, which asserts the existence of a matrix \tilde{S}_1 as in the first step in the above example earlier. (Note: The tilde indicates that this is not exactly the matrix S_1 mentioned in the example above, but this difference is explained and handled in Section 4.2.2.)

Proposition 4.2 (Constructing \tilde{S}_1). *Let $A_0 \in \mathbb{C}^{n \times n}$ be a fixed H_0 -selfadjoint matrix as given in (4.1). There exists a constant $K > 0$ (depending on A_0 and H_0 only) such that the following assertion holds. For any H -selfadjoint matrix A that is similar to $A_1 = J_n(\mu)$ for some μ there exists an affiliation matrix \tilde{S}_1 such that $(A, H) \xrightarrow{\tilde{S}_1} (A_1, \tilde{H}_1)$ for some lower anti-triangular*

Hankel matrix³ \tilde{H}_1 of the form $\begin{bmatrix} 0 & \cdots & 0 & * \\ \vdots & \ddots & \ddots & * \\ 0 & * & \ddots & \vdots \\ * & * & \cdots & * \end{bmatrix}$ satisfying

$$\|I - \tilde{S}_1\| \leq K \|A - A_0\|.$$

The proposition will be proved by the following lemma.

Lemma 4.3 (Bounds for the perturbed Jordan chains). *Let $A_0 \in \mathbb{C}^{n \times n}$ have only one eigenvalue λ . Let $\{f_k\}_{k=0}^{m-1}$ be the longest Jordan chain corresponding to the eigenvalue λ of A_0 . There are constants $K, \delta > 0$ (depending on A_0 only) such that for any $A \in \mathbb{C}^{n \times n}$ having only one eigenvalue μ and satisfying*

$$\|A - A_0\| \leq \delta,$$

we have the following. If the maximal length of the corresponding Jordan chain of A is also m , then there exists a Jordan chain $\{g_k\}_{k=0}^{m-1}$ of A such that

$$\|f_k - g_k\| \leq K \|A - A_0\|, \quad k = 0, \dots, m - 1.$$

This lemma is actually a specification of its more general version, Lemma 6.14, which will be proved later without using any intermediate results.

We are now ready to provide the proof of Proposition 4.2.

Proof of Proposition 4.2. Using the notations of Lemma 4.3, choose the matrix \tilde{S}_1 such that $\tilde{S}_1 f_i = g_i$, for $i = 0, \dots, n - 1$. We denote the result of applying the matrix \tilde{S}_1 to H by \tilde{H}_1 , so

$$(A, H) \xrightarrow{\tilde{S}_1} (A_1, \tilde{H}_1).$$

We next show that \tilde{H}_1 is lower anti-triangular Hankel. Writing $A_1 = \mu I + Z^T$ where Z denotes the lower shift matrix, and from the fact that A_1 is \tilde{H}_1 -selfadjoint we have that

³ By lower anti-triangular Hankel matrix we mean a matrix whose entries above the anti-diagonal are zeros.

$$\tilde{H}_1(\mu I + Z^T) = (\mu I + Z)\tilde{H}_1$$

and hence

$$\tilde{H}_1 Z^T = Z \tilde{H}_1$$

from which it immediately follows that \tilde{H}_1 is lower anti-triangular Hankel.

Next, for any $x \in \mathbb{C}^m$ with $\|x\| = 1$, write $x = \sum_{i=0}^{n-1} \alpha_i f_i$. Then $y = \tilde{S}_1 x = \sum_{i=0}^{n-1} \alpha_i g_i$, and

$$\|(I - \tilde{S}_1)x\| = \|x - y\| \leq \sum_{i=0}^{n-1} |\alpha_i| \|f_i - g_i\| \leq \max_{0 \leq i \leq n-1} |\alpha_i| \cdot \|f_i - g_i\|.$$

Next, denoting by $[x]_F = (\alpha_{i-1})$ the coordinates of x with respect to the fixed basis $\{f_i\}$ and $\mathcal{P}_{F \leftarrow E}$ the change of basis matrix from the standard basis $\{e_i\}$ to $\{f_i\}$, we have

$$\max_{0 \leq i \leq n-1} |\alpha_i| \leq \sqrt{\sum_{j=0}^{n-1} |\alpha_j|^2} = \|[x]_F\| \leq \|\mathcal{P}_{F \leftarrow E}\| \cdot \|x\| = \|\mathcal{P}_{F \leftarrow E}\|.$$

Using Lemma 4.3 we have that

$$\|(I - \tilde{S}_1)x\| \leq \|\mathcal{P}_{F \leftarrow E}\| \cdot \|f_i - g_i\| \leq K \|A - A_0\|$$

for any $x \in \mathbb{C}^m$ with $\|x\| = 1$. Hence

$$\|I - \tilde{S}_1\| \leq K \|A - A_0\| \tag{4.5}$$

as desired.

4.2.2. Modified first step. Forcing unit antidiagonal of H_1 . Constructing S_1

The matrix \tilde{S}_1 constructed in Section 4.2.1 was such that $(A, H) \xrightarrow{\tilde{S}_1} (A_1, \tilde{H}_1)$ with \tilde{H}_1 a lower anti-triangular Hankel matrix of the form

$$\tilde{H}_1 = \epsilon \cdot \begin{bmatrix} 0 & \cdots & 0 & a \\ \vdots & \ddots & a & * \\ 0 & \ddots & \ddots & \vdots \\ a & * & \cdots & * \end{bmatrix} \quad \text{with } \epsilon = \pm 1. \tag{4.6}$$

In the next proposition we construct a different matrix S_1 such that mapping $(A, H) \xrightarrow{S_1} (A_1, H_1)$ produces a better matrix H_1 of the form

$$H_1 = \epsilon \cdot \begin{bmatrix} 0 & \cdots & 0 & 1 \\ \vdots & \ddots & 1 & * \\ 0 & \ddots & \ddots & \vdots \\ 1 & * & \cdots & * \end{bmatrix} \quad \text{with } \epsilon = \pm 1 \tag{4.7}$$

(Recall that in the model example of this section the matrix H_1 of (4.4) indeed had the form (4.7)).

Proposition 4.4 (Modified Proposition 4.2. Constructing S_1). *Let $A_0 \in \mathbb{C}^{n \times n}$ be a fixed H_0 -selfadjoint matrix as given in (4.1). There exist constants $K, \delta > 0$ (depending on A_0 and H_0 only) such that the following assertion holds. For any H -selfadjoint matrix A that is similar to $A_1 = J_n(\mu)$ for some μ and*

$$\|A - A_0\| + \|H - H_0\| < \delta,$$

there exists an affiliation matrix S_1 such that

$$(A, H) \xrightarrow{S_1} (A_1, H_1) \text{ for some lower anti-triangular Hankel matrix } H_1 \text{ of the form (4.7)}$$

satisfying

$$\|I - S_1\| \leq K (\|A - A_0\| + \|H - H_0\|).$$

Let \tilde{S}_1 be the matrix guaranteed by Proposition 4.2, and a the antidiagonal entry of the resulting matrix \tilde{H}_1 as in (4.6). We will demonstrate that the matrix

$$S_1 := a^{-1/2} \tilde{S}_1$$

is the desired matrix. Indeed, it is easy to see that $S_1^{-1}AS_1 = A_1$ and $S_1^*HS_1 = H_1$ for H_1 of the form (4.7). The next lemma demonstrates that the bound on $\|I - \tilde{S}_1\|$ given by Proposition 4.2 implies the corresponding bound for $\|I - S_1\|$.

Lemma 4.5 (Forcing unit antidiagonal of G). *Let $A_0 \in \mathbb{C}^{n \times n}$ be an H_0 -selfadjoint matrix. There exist positive constants $K_1, K_2, \delta > 0$ (all depending on A_0 and H_0 only) such that the following assertion holds. For any H -selfadjoint matrix A satisfying*

$$\|A - A_0\| + \|H - H_0\| \leq \delta, \tag{4.8}$$

and any invertible matrix T satisfying

$$\|I - T\| \leq K_1(\|A - A_0\| + \|H - H_0\|) \tag{4.9}$$

and such that the matrix B in $(A, H) \xrightarrow{T} (B, G)$ is a single Jordan block we have

$$\|I - a^{-1/2}T\| \leq K_2(\|A - A_0\| + \|H - H_0\|), \tag{4.10}$$

where a is the $(n, 1)$ -entry of the matrix G .

Proof. The proof consists of two parts. In part (i) we prove that there is $K_3 > 0$ (depending on A_0 and H_0 only) such that

$$\begin{aligned} \|G - H_0\| &\leq K_3(\|A - A_0\| + \|H - H_0\|) \quad \text{with} \\ K_3 &= ((1 + K_1\delta)^2 + (2 + K_1\delta)\|H_0\|). \end{aligned} \tag{4.11}$$

Then we prove in part (ii) that the desired bound (4.10) is valid.

(i) (Proving (4.11)) We have

$$\begin{aligned} \|G - H_0\| &= \|T^*HT - H_0\| = \|T^*HT - T^*H_0T + T^*H_0T - T^*H_0 + T^*H_0 - H_0\| \\ &\leq \|T\|^2 \cdot \|H - H_0\| + (1 + \|T\|)(\|H_0\| \cdot \|T - I\|). \end{aligned} \tag{4.12}$$

From conditions (4.8) and (4.9), we have

$$\|T\| \leq 1 + K_1(\|A - A_0\| + \|H - H_0\|) \leq 1 + K_1\delta =: M \tag{4.13}$$

and substituting this estimate in (4.12) yields

$$\|G - H_0\| \leq \left[M^2 + (1 + M)\|H_0\| \right] (\|A - A_0\| + \|H - H_0\|),$$

which yields (4.11).

(ii) (Proving (4.10)) We first show that

$$|a - 1| \leq \tilde{K}_2(\|A - A_0\| + \|H - H_0\|) \quad \text{and} \quad a > \frac{1}{2} \tag{4.14}$$

for some constant \tilde{K}_2 . Using the relations

$$\epsilon = e_n^T H_0 e_1 \quad \text{and} \quad \epsilon \cdot a = e_n^T G e_1$$

with e_k the k th unit vector, and the fact that $\epsilon = \pm 1$, we have that

$$|a - 1| = |\epsilon \cdot a - \epsilon| = \|e_n^T (G - H_0) e_1\| \leq \|G - H_0\|$$

and the bound for $|a - 1|$ follows from (4.11). Next, since δ is at our disposal, we can always assume that it is small enough to guarantee that

$$\delta \leq (2\tilde{K}_2)^{-1},$$

so the condition (4.8) implies that

$$|a - 1| \leq \tilde{K}_2(\|A - A_0\| + \|H - H_0\|) < \tilde{K}_2 \delta \leq \frac{1}{2},$$

and hence $a > 1/2$, so both statements of (4.14) follow by perhaps considering a smaller neighborhood δ of the pair (A_0, H_0) .

Next,

$$\left\| I - \frac{1}{\sqrt{a}} T \right\| = \left\| I - T + T - \frac{1}{\sqrt{a}} T \right\| \leq \|I - T\| + \left| \frac{\sqrt{a} - 1}{\sqrt{a}} \right| \|T\|,$$

and using (4.14), we have that

$$\left| \frac{\sqrt{a} - 1}{\sqrt{a}} \right| = \left| \frac{a - 1}{a + \sqrt{a}} \right| \leq \left| \frac{a - 1}{a} \right| \leq 2|a - 1|.$$

Thus, we have that

$$\left\| I - \frac{1}{\sqrt{a}} T \right\| \leq \|I - T\| + 2|a - 1| \cdot \|T\|$$

and using the bounds for $\|T\|$ in (4.13) and $|a - 1|$ in (4.14), we arrive at

$$\left\| I - a^{-1/2} T \right\| \leq \left[K_1 + 2M(M^2 + (1 + M)\|H_0\|) \right] (\|A - A_0\| + \|H - H_0\|),$$

which establishes desired (4.10). \square

We next return to and prove Proposition 4.4.

Proof of Proposition 4.4. For the given pair (A_0, H_0) , Proposition 4.2 gives a constant K_1 such that for any applicable pair (A, H) there exists a matrix \tilde{S}_1 satisfying $(A, H) \xrightarrow{\tilde{S}_1} (A_1, \tilde{H}_1)$ for some lower anti-triangular Hankel matrix \tilde{H}_1 , and $\|I - \tilde{S}_1\| \leq K_1 \|A - A_0\|$. Applying Lemma 4.5 with $T = \tilde{S}_1$ and $G = \tilde{H}_1$, we have that the matrix $S_1 = a^{-1/2} \tilde{S}_1$, with $a = (\tilde{H}_1)_{n,1}$ satisfies the desired bound. Furthermore,

$$S_1^{-1} A S_1 = (a^{1/2} \tilde{S}_1^{-1}) A (a^{-1/2} \tilde{S}_1) = A_1$$

and

$$S_1^* H S_1 = (a^{-1/2} \tilde{S}_1^*) H (a^{-1/2} \tilde{S}_1) = a^{-1} \tilde{H}_1 = H_1$$

or

$$(A, H) \xrightarrow{S_1} (A_1, H_1)$$

by Proposition 4.2, where H_1 is then of the form (4.7) as desired. \square

4.2.3. Second step. Zeroing sub-antidiagonal entries of H_1 . Constructing S_2

With the modified first step, we are now in the same position as before the second step in the model example of this section. The next proposition implements the second step of the model example.

Proposition 4.6 (Zeroing sub-antidiagonal entries of H_1). *Let A_0 be a fixed H_0 -selfadjoint matrix as given in (4.1). Then there exist constants $K, \delta > 0$ (depending on A_0 and H_0 only) such that for any $A_1 = J_n(\mu)$ for some μ , and H_1 of the form shown in (4.7) satisfying*

$$\|A_1 - A_0\| + \|H_1 - H_0\| < \delta,$$

there exists an affiliation matrix S_2 such that

$$(A_1, H_1) \xrightarrow{S_2} (A_1, H_0)$$

satisfying

$$\|I - S_2\| \leq K (\|A_1 - A_0\| + \|H_1 - H_0\|).$$

The previous proposition guarantees the existence of a matrix S_2 that satisfies the following two properties:

- (1) $S_2^{-1} A_1 S_2 = A_1$, and
- (2) $S_2^* H_1 S_2 = H_0$.

That is, it must not affect the matrix A_1 , and it must zero out the sub-antidiagonal elements of the matrix H_1 . The former can be accomplished by choosing S_2 to be upper triangular Toeplitz, as since A_1 is also upper triangular Toeplitz, they will commute. For the latter, this is essentially accomplished by choosing a matrix that is the matrix square root of the inverse of the given Hankel matrix H_1 . The next lemma shows that this matrix has the desired properties.

Lemma 4.7. *Let $R \in \mathbb{R}^{n \times n}$ be a lower anti-triangular Hankel matrix with ones on the main antidiagonal. There is a constant $K > 0$ (depending on R only) such that the following statement holds. Denote by T the lower triangular Toeplitz matrix $T = R\tilde{I}$. Write $T = I + E$ and suppose $\|E\| \leq M$ for some bound $M > 1$.*

Define $S = (f(E))^*$, where

$$f(x) = \sqrt{\frac{1}{1+x}}.$$

Then

- (i) $\|I - S\| \leq K \|E\|$,

- (ii) S is upper triangular Toeplitz,
- (iii) $S^*RS = \tilde{I}$.

Proof. To prove (i) note that it follows from the definitions that E is nilpotent with index of nilpotence at most n . Thus we have $E^n = 0$, and using the power series expansion for $\sqrt{\frac{1}{1+x}}$ we see that

$$S = f(E) = \sum_{k=0}^{n-1} \left(\prod_{j=1}^k \frac{1-2j}{2j} \right) E^k$$

and so

$$\|I - S\| \leq \sum_{k=1}^{n-1} \|E\|^k \leq \left(\sum_{k=1}^{n-1} M^{k-1} \right) \|E\| \leq (n - 1)M^{n-2} \|E\|.$$

To prove (ii) note that from (i) S is defined as a transpose of a finite linear combination of powers of a lower triangular Toeplitz matrix E , so S is an upper triangular Toeplitz matrix.

(iii) From the definition of S , $I = S^*S^*T$, and since S^* and T are both lower triangular Toeplitz matrices, S^* and T commute hence

$$I = S^*TS^* = S^*R\tilde{I}S^*.$$

Postmultiplying both side with \tilde{I} gives that

$$\tilde{I} = S^*R\tilde{I}S^*\tilde{I} = S^*RS,$$

completing the proof. \square

We are now ready to return to the proof of Proposition 4.6.

Proof of Proposition 4.6. Letting $R = \epsilon H_1$ where ϵ is the sign characteristic of the pair (A_0, H_0) and applying Lemma 4.7 (the fact that $\|E\| \leq M$ for some M follows from (4.11), proved in the proof of Lemma 4.5) gives a matrix S_2 such that $S_2^*(\epsilon H_1)S_2 = \tilde{I}$, and hence $S_2^*H_1S_2 = H_0$. Since S_2 is an upper triangular Toeplitz matrix, it commutes with A_1 which is also an upper triangular Toeplitz matrix, and so $S_2^{-1}A_1S_2 = A_1$. These two facts together yield

$$(A_1, H_1) \xrightarrow{S_2} (A_1, H_0).$$

To prove the bound on $\|I - S_2\|$, note that with $R = \epsilon H_1$ and the notations of Lemma 4.7, we have that $\epsilon H_1\tilde{I} = I + E$ which implies $E = \epsilon H_1\tilde{I} - I$. On using the fact that $\|\epsilon\tilde{I}\| = 1$ we have that

$$\|E\| = \|\epsilon E\tilde{I}\| = \|H_1 - H_0\| \leq K(\|A - A_0\| + \|H - H_0\|)$$

from (4.11) of the proof of Lemma 4.5.

4.2.4. Third step. Combining affiliation matrices S_1 and S_2

We have seen in step 3 of the model example of Section 4.1 that we can combine the affiliation matrices S_1 and S_2 produced by the previous steps to obtain the desired matrix. That is, we already have that

$$(A, H) \xrightarrow{S_1} (A_1, H_1) \xrightarrow{S_2} (A_1, H_0)$$

and hence the matrix $S = S_1 S_2$ is such that

$$(A, H) \xrightarrow{S} (A_1, H_0).$$

Moreover, individual Lipschitz-type bounds for S_1 and S_2 yielded an overall bound of the same form for S . In order to accomplish this in general and combine the matrices S_1 and S_2 of Propositions 4.4 and 4.6, respectively, we will need the following auxiliary result.

Lemma 4.8 (Near-identity affiliation matrix yields small perturbations). *Let $A_0 \in \mathbb{C}^{n \times n}$ be an H_0 -selfadjoint matrix, and A be an H -selfadjoint matrix that has the same Jordan structure as A_0 , and T be an invertible matrix satisfying $(A, H) \xrightarrow{T} (B, G)$ for some G -selfadjoint matrix B . Suppose that there exist constants $K, \delta > 0$ such that*

$$\|I - T\| \leq K(\|A - A_0\| + \|H - H_0\|)$$

and

$$\|A - A_0\| + \|H - H_0\| < \delta.$$

Then

$$\|G - H_0\| \leq K_1(\|A - A_0\| + \|H - H_0\|) \quad \text{with} \quad K_1 = (1 + K\delta)^2 + (2 + K\delta)\|H_0\|,$$

and

$$\|B - A_0\| \leq K_2(\|A - A_0\| + \|H - H_0\|) \quad \text{with} \quad K_2 = 2K(1 + K\delta) + 4K\|A_0\|.$$

Proof. The bound for $\|G - H_0\|$ is the same as the bound (4.11), and was established in the first part of the proof of Lemma 4.5. To prove the second bound, we have (similar to the proof for $\|G - H_0\|$) that

$$\|B - A_0\| \leq \|T^{-1}\| \cdot \|T\| \cdot \|A - A_0\| + 2\|A_0\| \cdot \|T^{-1}\| \cdot \|I - T\|$$

by using the obvious identity $\|I - T^{-1}\| \leq \|T^{-1}\| \cdot \|I - T\|$. Next, by perhaps considering a smaller δ , we can assume that δ is small enough so that $\|I - T\| \leq \frac{1}{2}$. Then

$$\|T^{-1}\| - 1 \leq \|I - T^{-1}\| \leq \|T^{-1}\| \cdot \|I - T\| \leq \frac{1}{2}\|T^{-1}\|,$$

which leads to

$$\|T^{-1}\| \leq 2.$$

Using this bound for $\|T^{-1}\|$ in combination with the bound for $\|T\| \leq M$ of (4.13) in Lemma 4.5, we have

$$\|B - A_0\| \leq [2KM + 4K\|A_0\|](\|A - A_0\| + \|H - H_0\|). \quad \square$$

Theorem 4.9 (Combining affiliation matrices S_1 and S_2). *Let $A_0 \in \mathbb{C}^{n \times n}$ be a fixed H_0 -selfadjoint matrix. There exist positive constants K, δ (all depending on A_0 and H_0 only) such that for any H -selfadjoint matrix A satisfying*

$$\|A - A_0\| + \|H - H_0\| < \delta, \tag{4.15}$$

the following statement is true. If we consider mappings

$$(A, H) \xrightarrow{S_1} (B, G) \xrightarrow{S_2} (C, F)$$

with any S_1, S_2 satisfying

$$\|I - S_1\| \leq K(\|A - A_0\| + \|H - H_0\|), \quad \|I - S_2\| \leq K(\|B - A_0\| + \|G - H_0\|), \tag{4.16}$$

then $S = S_1 S_2$ satisfies $(A, H) \xrightarrow{S} (C, F)$ and

$$\|I - S\| \leq (2K + K^2\delta)(\|A - A_0\| + \|H - H_0\|).$$

Proof. We compute

$$S^{-1}AS = S_2^{-1}S_1^{-1}AS_1S_2 = S_2^{-1}BS_2 = C$$

and

$$S^*HS = S_2^*S_1^*HS_1S_2 = S_2^*GS_2 = F$$

using $(A, H) \xrightarrow{S_1} (B, G)$ and then $(B, G) \xrightarrow{S_2} (C, F)$ in each computation. This establishes $(A, H) \xrightarrow{S} (C, F)$. From (4.16) and (4.15),

$$\|S_i\| \leq 1 + K\delta, \quad i = 1, 2.$$

Therefore, the matrix $S = S_1 S_2$ satisfies

$$\begin{aligned} \|I - S\| &\leq \|I - S_1\| + \|S_1\| \cdot \|I - S_2\| \leq \|I - S_1\| + (1 + K\delta) \cdot \|I - S_2\| \\ &\leq (2K + K^2\delta)(\|A - A_0\| + \|H - H_0\|) \end{aligned}$$

as claimed. \square

Proof of Theorem 4.1 in the case of a real single Jordan block. From Lemma 4.8 we have the individual bounds

$$\|I - S_i\| \leq K_i(\|A - A_0\| + \|H - H_0\|), \quad i = 1, 2.$$

Hence applying Theorem 4.9 yields exactly (4.3), completing the justification of Theorem 4.1 in the case of (A_0, H_0) in the real single Jordan block case. \square

In the next section, this result is expanded to the single complex eigenvalue case.

4.3. Sketch of the proof of Theorem 4.1 in the case of two complex conjugate Jordan blocks

In Section 4.2, Theorem 4.1 was completely proved for the single *real* Jordan block case. The results need to be modified slightly to complete the proof of Theorem 4.1 and to prove it for the case of a single complex eigenvalue pair.⁴ Specifically, we now consider the case where

$$A_0 = \begin{bmatrix} J(\lambda) & 0 \\ 0 & J(\bar{\lambda}) \end{bmatrix}, \quad H_0 = \tilde{I}, \tag{4.17}$$

where we use the notations of (2.1).

⁴ Recall that the eigenvalues of H -selfadjoint matrices are either real or occur in complex conjugate pairs having identical Jordan structure, see, e.g., Theorem 2.1.

For some pair (A, H) with the same Jordan structure as (A_0, H_0) (that is, A has Jordan form $A_1 = J_k(\mu) \oplus J_k(\bar{\mu})$ for some complex μ), suppose that a matrix S_1 is found as in Section 4.2.1 such that

$$(A, H) \xrightarrow{S_1} (A_1, H_1).$$

We demonstrate that the matrix H_1 must have the form

$$H_1 = \begin{bmatrix} 0 & G^* \\ G & 0 \end{bmatrix}, \quad G = (g_{i+j}) = \begin{bmatrix} 0 & \cdots & 0 & g_{n+1} \\ \vdots & \ddots & \ddots & g_{n+2} \\ 0 & \ddots & \ddots & \vdots \\ g_{n+1} & g_{n+2} & \cdots & g_{2n} \end{bmatrix}. \tag{4.18}$$

Indeed, from [6, Corollary 4.2.5], it follows that the upper-left and lower-right blocks of H_1 are all zeros as claimed. Further, A_1 is H_1 -selfadjoint, and so $H_1 A_1$ must be selfadjoint and

$$H_1 A_1 = \begin{bmatrix} 0 & G^* J(\bar{\mu}) \\ G J(\mu) & 0 \end{bmatrix},$$

hence

$$G J(\mu) = (G^* J(\bar{\mu}))^*$$

and denoting by Z the lower shift matrix of appropriate size,

$$G Z^T = Z G$$

and so G is a Hankel matrix as claimed. From this point, Proposition 4.6 can be suitably modified to produce a matrix S_2 such that

$$(A, H) \xrightarrow{S_1} (A_0, H_1) \xrightarrow{S_2} (A_0, H_0)$$

and hence the matrix $S = S_1 S_2$ is as desired.

Theorem 4.1 is now completely proved.

5. Proof of Theorem 1.8 for the multiple Jordan block case

In Section 4 the main result, Theorem 1.8, was proved for the case where (A_0, H_0) are in the canonical form of Theorem 2.1 and A_0 was either a single real Jordan block or a direct sum of two conjugate nonreal Jordan blocks. The following theorem generalizes this result to the case where A_0 is an arbitrary Jordan canonical form matrix. In accordance with the Theorem 3.1, this will completely prove the desired Theorem 1.8.

Theorem 5.1 (Extension of Theorem 4.1. Lipschitz stability of affiliation matrices in the multi-block case). *Let $A_0 \in \mathbb{C}^{n \times n}$ be a fixed H_0 -selfadjoint matrix, both A_0 and H_0 in the canonical form described in Theorem 2.1. Then there exist constants $K, \delta > 0$ (depending on A_0 and H_0 only) such that the following assertion holds. For any H -selfadjoint matrix A such that A has the same Jordan structure as A_0 and*

$$\|A - A_0\| + \|H - H_0\| < \delta,$$

the pairs (A_0, H_0) and (A, H) are affiliated, and there exists a affiliation matrix S such that

$$\|I - S\| \leq K(\|A - A_0\| + \|H - H_0\|). \tag{5.1}$$

This theorem will be proved by induction on the number of the Jordan blocks of A_0 . Theorem 4.1 of Section 4 establishes the result for the case when A_0 has a single block $\tilde{J}(\lambda)$. Now, to make the inductive step we will need the following result that allows us to “decouple” such individual blocks from the rest.

Lemma 5.2 (Decoupling). *Let $A_0 \in \mathbb{C}^{n \times n}$ be a fixed H_0 -selfadjoint matrix, both A_0 and H_0 in the canonical form described in Theorem 2.1, and denote*

$$A_0 = \begin{bmatrix} \tilde{J}(\lambda_1) & 0 \\ 0 & \tilde{A}_0 \end{bmatrix} \quad \text{and} \quad H_0 = \begin{bmatrix} \epsilon_1 P_1 & 0 \\ 0 & \tilde{H}_0 \end{bmatrix}; \tag{5.2}$$

i.e. consider the partition that singles out a block $\tilde{J}(\lambda_1)$ of the form defined in (2.1). We assume that $\tilde{J}(\lambda_1)$ is the biggest block of A_0 corresponding to the eigenvalue λ_1 . Then there are positive constants K and δ (depending on A_0 and H_0 only) such that for any H -selfadjoint matrix A with the same Jordan structure as A_0 satisfying

$$\|A - A_0\| + \|H - H_0\| < \delta,$$

the following statements hold.

(i) *A has an eigenvalue μ_1 satisfying*

$$|\lambda_1 - \mu_1| \leq K \|A - A_0\|. \tag{5.3}$$

(ii) *There exists an affiliation matrix S , i.e., $(A, H) \xrightarrow{S} (A_2, H_2)$ such that the matrices A_2 and H_2 have the form*

$$A_2 = \begin{bmatrix} \tilde{J}(\mu_1) & 0 \\ 0 & \tilde{A}_1 \end{bmatrix} \quad \text{and} \quad H_2 = \begin{bmatrix} \epsilon_1 P_1 & 0 \\ 0 & \tilde{H}_1 \end{bmatrix}, \tag{5.4}$$

with some \tilde{A}_1, \tilde{H}_1 and S satisfies the bound

$$\|I - S\| \leq K (\|A - A_0\| + \|H - H_0\|). \tag{5.5}$$

The proof of this lemma is postponed to Section 7. Lemma 5.2 allows us to make an inductive step and to prove Theorem 5.1.

Proof of Theorem 5.1. Applying Lemma 5.2 to the pair (A, H) , as in

$$(A, H) \xrightarrow{S_1} (A_1, H_1),$$

results in a pair (A_1, H_1) of the form shown in (5.4) with

$$\|I - S_1\| \leq K_1 (\|A - A_0\| + \|H - H_0\|). \tag{5.6}$$

Since the lower right block \tilde{A}_1 of A_1 has less Jordan blocks than A_0 , we can use the inductive assumption and apply Theorem 5.1 to the pair $(\tilde{A}_1, \tilde{H}_1)$, as in

$$(\tilde{A}_1, \tilde{H}_1) \xrightarrow{\tilde{S}_2} (\tilde{A}_2, \tilde{H}_2),$$

with

$$\|I - \tilde{S}_2\| \leq K_2 (\|\tilde{A}_1 - \tilde{A}_0\| + \|\tilde{H}_1 - \tilde{H}_0\|). \tag{5.7}$$

From this follows

$$\|I - S_2\| \leq K'_2(\|A_2 - A_0\| + \|H_2 - H_0\|) \tag{5.8}$$

with

$$S_2 = \begin{bmatrix} I & 0 \\ 0 & \tilde{S}_2 \end{bmatrix}.$$

Indeed, the transition from (5.7) to (5.8) involves only adding the (1, 1) blocks into consideration, and from (5.2) and (5.4), we see that the only nontrivial addition is from $\|J(\mu_1) - J(\lambda_1)\|$. In view of Lemma 6.14 (which is proved independently of results of this theorem), this serves only to modify the constant, and (5.8) is established. Observe, that from (6.33) and (5.7) it follows that $K'_2 = \max\{1, K_2\}$ depends on A_0 and H_0 only.

Finally, Theorem 4.9 allows us to combine the bounds (5.6) and (5.8), and it implies the desired bound (5.5) for $S = S_1 S_2$. \square

To complete the proof of Theorem 5.1 (and thus, of the main result, Theorem 1.8), it remains only to prove Lemma 5.2. It will be done in Section 7 after we obtain in the next auxiliary section several necessary results on the perturbation of subspaces.

6. Auxiliary lemmas on semigaps, gaps and perturbations of subspaces

To provide the proof of Lemma 5.2 in Section 7 one needs to use a number of results on small perturbations of certain invariant subspaces. We have gathered all these auxiliary results in this section. In order to obtain specific bounds for such perturbations, we need to deal with the distance between two subspaces. One standard way to define such a distance is based on the concept of a *gap*. Our approach below is slightly different, it is based on the concept of a *semigap* for which it is often easier to obtain the desired bounds. The key result of this section is Lemma 6.6 that says that when two subspaces have equal dimensions the gap between them is equal to the semigap. Many of the auxiliary results in this section are known. For somewhat less known results, e.g., about semigaps, we provide references and in some cases new proofs.

6.1. Gap between subspaces

Before defining the concept of a gap, let us recall that

- A matrix $P_{\mathcal{M}}$ is called a *projector* onto a subspace $\mathcal{M} \subset \mathbb{C}^n$ if (a) $\text{Im } P_{\mathcal{M}} = \mathcal{M}$; (b) $P_{\mathcal{M}}^2 = P_{\mathcal{M}}$.
- Further, $P_{\mathcal{M}}$ is called an *orthogonal projector* onto \mathcal{M} if additionally we have (c) $P_{\mathcal{M}}^* = P_{\mathcal{M}}$.

Here is the key definition.

Definition 6.1 (*Gap. First definition*). The gap $\theta(\mathcal{M}, \mathcal{N})$ between two subspaces $\mathcal{M}, \mathcal{N} \subset \mathbb{C}^n$ can be introduced via

$$\theta(\mathcal{M}, \mathcal{N}) = \|P_{\mathcal{M}} - P_{\mathcal{N}}\|, \tag{6.1}$$

where $P_{\mathcal{M}}$ denotes the orthogonal projector onto \mathcal{M} .

It is well-known, see, e.g., [5], that *gap* is a metric in the set of all subspaces of \mathbb{C}^n .

The definition in (6.1) has been found to useful in many instances. However, in Theorem 2.6 and Lemma 4.3 above we did not deal with entire subspaces, but rather with particular vectors

spanning them. Therefore, in our context, it is often more convenient to use the next definition that is well-known to be equivalent to the first one.

Definition 6.2 (*Gap. Second definition*). The gap $\theta(\mathcal{M}, \mathcal{N})$ between two subspaces $\mathcal{M}, \mathcal{N} \subset \mathbb{C}^n$ can be introduced via

$$\theta(\mathcal{M}, \mathcal{N}) = \max \left\{ \sup_{\substack{x \in \mathcal{M} \\ \|x\|=1}} \inf_{y \in \mathcal{N}} \|x - y\|, \sup_{\substack{y \in \mathcal{N} \\ \|y\|=1}} \inf_{x \in \mathcal{M}} \|y - x\| \right\}. \tag{6.2}$$

Theorem 2.6 and Lemma 4.3 have just been used as a motivation for (6.2), but in these statements the vectors $\{f_k\}$ were fixed, while the vectors $\{g_k\}$ were their perturbations. One might expect that in some instances it might be easier to rely on the properties of fixed vectors $\{f_k\}$, but the quantity $\theta(\text{span}\{f_k\}, \text{span}\{g_k\})$ is clearly symmetric, so it does not give $\{f_k\}$'s any "advantage." In order to better capture the difference between $\{f_k\}$'s and $\{g_k\}$'s we now introduce a different one-sided quantity.

6.2. Semigap between subspaces

Following [14] we give the following definition.

Definition 6.3 (*Semigap*). Let $\mathcal{M}, \mathcal{N} \subset \mathbb{C}^n$ be two subspaces. The quantity

$$\theta_0(\mathcal{M}, \mathcal{N}) = \sup_{\substack{x \in \mathcal{M} \\ \|x\|=1}} \inf_{y \in \mathcal{N}} \|x - y\|$$

is called the semigap (or one-sided gap) from \mathcal{M} to \mathcal{N} .

Clearly, in light of Definition 6.2,

$$\theta(\mathcal{M}, \mathcal{N}) = \max\{\theta_0(\mathcal{M}, \mathcal{N}), \theta_0(\mathcal{N}, \mathcal{M})\}.$$

Lemma 6.4. [Some immediate properties of semigaps] For any two subspaces $\mathcal{M}, \mathcal{N} \subset \mathbb{C}^n$ we have

$$\theta_0(\mathcal{M}, \mathcal{N}) = \sup_{\substack{x \in \mathcal{M} \\ \|x\|=1}} \|x - P_{\mathcal{N}}x\|, \tag{6.3a}$$

$$\text{If } \mathcal{N}_1 \subset \mathcal{N}_2, \text{ then } \theta_0(\mathcal{M}, \mathcal{N}_2) \leq \theta_0(\mathcal{M}, \mathcal{N}_1), \quad \theta_0(\mathcal{N}_1, \mathcal{M}) \leq \theta_0(\mathcal{N}_2, \mathcal{M}) \tag{6.3b}$$

$$\theta_0(\mathcal{M}, \mathcal{N}) \leq 1, \tag{6.3c}$$

$$\text{If } \dim \mathcal{M} > \dim \mathcal{N}, \text{ then } \theta_0(\mathcal{M}, \mathcal{N}) = 1. \tag{6.3d}$$

The four auxiliary assertions of this lemma are well known, and can be found, e.g., in [8,5].

The next example shows that generally the two quantities $\theta_0(\mathcal{M}, \mathcal{N})$ and $\theta_0(\mathcal{N}, \mathcal{M})$ need not coincide.

Example 6.5 (*Semigaps can be nonsymmetric*). Let $\mathcal{N} = \mathbb{C}^2$ and $\mathcal{M} = \text{span}\{e_1\}$. Since in this case $\mathcal{M} \subset \mathcal{N}$ hence

$$\theta_0(\mathcal{M}, \mathcal{N}) = 0.$$

However, the vector $e_2 \in \mathcal{N}$ is orthogonal to \mathcal{M} and hence

$$\theta_0(\mathcal{N}, \mathcal{M}) = 1.$$

In the above example the dimensions of \mathcal{M} and \mathcal{N} were different. The next statement shows that when the dimensions of \mathcal{M} and \mathcal{N} are the same then the two associated semigaps are always equal.

Lemma 6.6 (Semigaps for subspaces of the same dimension). *Let $\mathcal{M}, \mathcal{N} \subset \mathbb{C}^n$ be two arbitrary subspaces. If $\dim \mathcal{M} = \dim \mathcal{N}$, then*

$$\theta_0(\mathcal{M}, \mathcal{N}) = \theta_0(\mathcal{N}, \mathcal{M}) \quad (= \theta(\mathcal{M}, \mathcal{N})). \tag{6.4}$$

Proof. In order to prove (6.4) we need to consider three cases.

Case 1. $\dim \mathcal{M} = \dim \mathcal{N} = 1$. If $\mathcal{M} = \mathcal{N}$ there is nothing to prove, so we assume $\mathcal{M} \neq \mathcal{N}$.

- *Defining an appropriate (complex) Householder reflection.* Let us choose $x \in \mathcal{N}$ and $y \in \mathcal{M}$ such that $\|x\| = \|y\| = 1$. By an appropriate unimodular rescaling of y one can guarantee that $x^*y \in \mathbb{R}$. Further, let us define

$$U = I - 2ww^*, \quad \text{where } w = \frac{1}{\|x - y\|}(x - y).$$

It is well known (and can be easily checked) that the Householder reflection U is Hermitian and unitary. In particular,

$$U^2 = I. \tag{6.5}$$

Secondly, it can be easily checked that

$$Ux = y, \quad Uy = x. \tag{6.6}$$

Indeed,

$$Ux = x - 2 \frac{(y - x)(y - x)^*}{(y - x)^*(y - x)}x = x - \frac{(x - y)(x^*y - 1)}{y^*x - 1} = y.$$

The second equation in (6.6) follows from the first one and (6.5).

Finally, observe

$$UP_{\mathcal{N}}U = P_{\mathcal{M}}. \tag{6.7}$$

- *Proving $\theta_0(\mathcal{M}, \mathcal{N}) = \theta_0(\mathcal{N}, \mathcal{M})$.* From (6.6) and (6.7) and from the fact that U is unitary it follows that

$$\|x - P_{\mathcal{N}}x\| = \|Uy - UUP_{\mathcal{N}}Uy\| = \|U(y - P_{\mathcal{M}}y)\| = \|y - P_{\mathcal{M}}y\|.$$

This and the property (6.3a) imply the desired (6.4).

Case 2. $\dim \mathcal{M} = \dim \mathcal{N} = k > 1$ and $\theta_0(\mathcal{M}, \mathcal{N}) = 1$. First we observe that

$$\theta_0(\mathcal{M}, \mathcal{N}) = 1 \iff \mathcal{M} \cap \mathcal{N}^\perp \neq \{0\} \tag{6.8}$$

and

$$\theta_0(\mathcal{N}, \mathcal{M}) = 1 \iff \mathcal{N} \cap \mathcal{M}^\perp \neq \{0\} \tag{6.9}$$

and our task is to show that (6.8) implies (6.9).

Denote by \mathcal{M}_1 the orthogonal complement to $\mathcal{M} \cap \mathcal{N}^\perp$ in \mathcal{M} and define $\mathcal{N}_1 = P_{\mathcal{N}}\mathcal{M}_1$. In view of (6.8) we have $\dim \mathcal{M}_1 < \dim \mathcal{M}$ and hence $\dim \mathcal{N}_1 < \dim \mathcal{N}$. Therefore there exist $y \in \mathcal{N}$ that is orthogonal to \mathcal{N}_1 . Clearly, this y is orthogonal to \mathcal{M} implying (6.9).

Case 3. $\dim \mathcal{M} = \dim \mathcal{N} = k > 1$ and $\theta_0(\mathcal{M}, \mathcal{N}) < 1$.

In order to prove (6.4) we first observe that

$$P_{\mathcal{M}}\mathcal{N} = \mathcal{M}, \quad \text{and} \quad P_{\mathcal{N}}\mathcal{M} = \mathcal{N}. \tag{6.10}$$

Indeed, the result proved in case 2 above implies that if $\theta_0(\mathcal{M}, \mathcal{N}) < 1$ we must also have $\theta_0(\mathcal{N}, \mathcal{M}) < 1$. Therefore none of \mathcal{M}, \mathcal{N} contains vectors orthogonal to each other and (6.10) follows.

Now, since the unit circle in \mathcal{M} is compact hence the supremum in (6.3a) is attained, i.e., there exists $x \in \mathcal{M}$ such that $\|x\| = 1$, we have that

$$\theta_0(\mathcal{M}, \mathcal{N}) = \|x - P_{\mathcal{N}}x\|.$$

Denoting $\mathcal{M}_x = \text{span}\{x\}$ we have

$$\theta_0(\mathcal{M}_x, \mathcal{N}) = \theta_0(\mathcal{M}, \mathcal{N}). \tag{6.11}$$

Now, in view of (6.10) there is a subspace $\mathcal{N}_x \subset \mathcal{N}$ such that

$$\mathcal{M}_x = P_{\mathcal{M}}\mathcal{N}_x.$$

Due to this particular choice of \mathcal{N}_x we have

$$\theta_0(\mathcal{N}_x, \mathcal{M}_x) = \theta_0(\mathcal{N}_x, \mathcal{M}). \tag{6.12}$$

Since both \mathcal{M}_x and \mathcal{N}_x are one-dimensional, we have

$$\theta_0(\mathcal{M}_x, \mathcal{N}_x) = \theta_0(\mathcal{N}_x, \mathcal{M}_x). \tag{6.13}$$

Gathering all the above we have

$$\begin{aligned} \theta_0(\mathcal{M}, \mathcal{N}) &\stackrel{(6.11)}{=} \theta_0(\mathcal{M}_x, \mathcal{N}) \stackrel{(6.3b)}{\leq} \theta_0(\mathcal{M}_x, \mathcal{N}_x) \stackrel{(6.13)}{=} \theta_0(\mathcal{N}_x, \mathcal{M}_x) \stackrel{(6.12)}{=} \theta_0(\mathcal{N}_x, \mathcal{M}) \\ &\leq (6.3b)\theta_0(\mathcal{N}, \mathcal{M}) \end{aligned}$$

We have proved that $\theta_0(\mathcal{M}, \mathcal{N}) \leq \theta_0(\mathcal{N}, \mathcal{M})$ without making any assumptions on \mathcal{M} and \mathcal{N} . Hence the desired result (6.4) follows by symmetry.

The proof is complete. \square

Corollary 6.7. *If $\dim \mathcal{M} = \dim \mathcal{N}$ then*

$$\theta(\mathcal{M}, \mathcal{N}) = \sup_{\substack{x \in \mathcal{M} \\ \|x\|=1}} \|x - P_{\mathcal{N}}x\| = \sup_{\substack{x \in \mathcal{M} \\ \|x\|=1}} \inf_{y \in \mathcal{N}} \|x - y\|.$$

In [14] the above lemma and corollary were found to be useful to study the change of Jordan structure of H -selfadjoint matrices under small perturbations.

Lemma 6.8. *Let $\mathcal{M} = \text{span}\{f_i\}_{i=0}^{m-1}$ where vectors $\{f_i\}$ are linearly independent. There exists a constant $K > 0$ (depending on $\{f_i\}$ only) such that for any set of vectors $\{g_i\}_{i=0}^{m-1}$ satisfying*

$$\|f_i - g_i\| \leq K,$$

we have

$$\theta_0(\mathcal{M}, \mathcal{N}) \leq mK \|\mathcal{P}_{F \leftarrow E}\|, \tag{6.14}$$

where $\mathcal{N} = \text{span}\{g_i\}$, and $\mathcal{P}_{F \leftarrow E}$ denotes the change-of-coordinates matrix from the standard basis $E = \{e_i\}$ to $F = \{f_i\}$.

Proof. Let $x \in \mathcal{M}$ and $\|x\| = 1$. For the decomposition of x with respect to the fixed basis $F\{f_i\}$:

$$x = \alpha_0 f_0 + \dots + \alpha_{m-1} f_{m-1},$$

let us consider

$$y = \alpha_0 g_0 + \dots + \alpha_{m-1} g_{m-1}.$$

Clearly, $y \in \mathcal{N}$ and

$$\|x - y\| \leq \alpha_0 \|f_0 - g_0\| + \dots + \alpha_{m-1} \|f_{m-1} - g_{m-1}\| \leq \max_{0 \leq k \leq m-1} |\alpha_k| \cdot mK. \tag{6.15}$$

In order to complete the proof we need to find a bound on $\max_{0 \leq k \leq m-1} |\alpha_k|$. To this end let us consider two decompositions of x with respect to the fixed basis $F = \{f_i\}$ and the standard (also fixed) basis $E = \{e_i\}$, respectively:

$$x = \alpha_1 f_1 + \dots + \alpha_n f_n = x_1 e_1 + \dots + x_n e_n,$$

i.e.,

$$x = [x]_E = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}, \quad [x]_F = \begin{bmatrix} \alpha_1 \\ \alpha_2 \\ \vdots \\ \alpha_n \end{bmatrix}.$$

Now,

$$\max_{0 \leq k \leq m-1} |\alpha_k| \leq \sqrt{|\alpha_0|^2 + \dots + |\alpha_{m-1}|^2} = \|[x]_F\| \leq \|\mathcal{P}_{F \leftarrow E} \cdot [x]_E\| \leq \|\mathcal{P}_{F \leftarrow E}\| \tag{6.16}$$

and we see that (6.16) and (6.15) imply (6.14). \square

6.3. Bounds for the perturbations of H -orthogonal companions

The following definition [4] introduces a counterpart $\mathcal{M}^{[\perp]H}$ (or simply $\mathcal{M}^{[\perp]}$) of the usual orthogonal complement \mathcal{M}^\perp for the spaces with indefinite inner product induced by H .

Definition 6.9 (H -orthogonal companion). Let $H \in \mathbb{C}^n$ be an invertible Hermitian matrix. For a subspace $\mathcal{M} \subset \mathbb{C}^n$ its H -orthogonal companion is defined as

$$\mathcal{M}^{[\perp]} = \{x \in \mathbb{C}^n : [x, y]_H = 0 \forall y \in \mathcal{M}\}.$$

In Section 7 we will need to develop an inductive decoupling process passing from a given subspace \mathcal{M} and its small perturbation \mathcal{N} to their orthogonal companions $\mathcal{M}^{[\perp]}$ and $\mathcal{N}^{[\perp]}$, respectively. The following lemma will be key in this framework since it claims that in this case the subspace $\mathcal{M}^{[\perp]}$ is a small perturbation of $\mathcal{N}^{[\perp]}$.

Lemma 6.10 (Bounds for H -orthogonal companions). Let H_0 be a fixed invertible Hermitian matrix, and let $\mathcal{M} \subset \mathbb{C}^n$ be a fixed subspace. There is a constant $\delta > 0$ (depending on H_0 and \mathcal{M} only) such that for any invertible Hermitian H satisfying

$$\|H - H_0\| \leq \delta,$$

and any subspace \mathcal{N} satisfying

$$\dim \mathcal{M} = \dim \mathcal{N}, \quad \theta_0(\mathcal{M}, \mathcal{N}) \leq L \tag{6.17}$$

(with certain L) we have

$$\theta(\mathcal{M}^{[\perp]}, \mathcal{N}^{[\perp]}) \leq (\kappa(H_0) + 2\|H_0\|) \cdot (L + \|H - H_0\|). \tag{6.18}$$

Proof. First observe that $H^{-1}P_{\mathcal{M}^\perp}H$ is a projection onto subspace $H^{-1}\mathcal{M}^\perp$, though not necessarily orthogonal. Second, it is well known [6] that for two (possibly not orthogonal) projectors $Q_{\mathcal{M}}$ and $Q_{\mathcal{N}}$ onto \mathcal{M} and \mathcal{N} , respectively, we have

$$\|P_{\mathcal{M}} - P_{\mathcal{N}}\| \leq \|Q_{\mathcal{M}} - Q_{\mathcal{N}}\|.$$

Third, it is well known [6] that

$$\mathcal{M}^{[\perp]} = H_0^{-1}\mathcal{M}^\perp, \quad \mathcal{N}^{[\perp]} = H^{-1}\mathcal{N}^\perp.$$

From the above three facts it follows that

$$\theta(\mathcal{M}^{[\perp]}, \mathcal{N}^{[\perp]}) = \|P_{H_0^{-1}\mathcal{M}^\perp} - P_{H^{-1}\mathcal{N}^\perp}\| \leq \|H_0^{-1}P_{\mathcal{M}^\perp}H_0 - H^{-1}P_{\mathcal{N}^\perp}H\|. \tag{6.19}$$

To bound the latter let us proceed with the right hand side of (6.19), and add and subtract the quantity $H_0^{-1}P_{\mathcal{N}^\perp}H_0$ and $H_0^{-1}P_{\mathcal{N}^\perp}H$:

$$\begin{aligned} & \| (H_0^{-1}P_{\mathcal{M}^\perp}H_0 - H_0^{-1}P_{\mathcal{N}^\perp}H_0) + (H_0^{-1}P_{\mathcal{N}^\perp}H_0 - H_0^{-1}P_{\mathcal{N}^\perp}H) \\ & \quad + (H_0^{-1}P_{\mathcal{N}^\perp}H - H^{-1}P_{\mathcal{N}^\perp}H) \| \\ & \leq \kappa(H_0)\|P_{\mathcal{M}^\perp} - P_{\mathcal{N}^\perp}\| + \|H_0^{-1}\| \cdot \|H - H_0\| + \|H_0^{-1}\| \cdot \|H - H_0\|, \end{aligned} \tag{6.20}$$

where $\kappa(H_0) = \|H_0\| \cdot \|H_0^{-1}\|$. Combining (6.19) and (6.20) we finally obtain

$$\theta(\mathcal{M}^{[\perp]}, \mathcal{N}^{[\perp]}) \leq \kappa(H_0) \cdot \theta(\mathcal{M}^\perp, \mathcal{N}^\perp) + 2\|H_0^{-1}\| \cdot \|H_0 - H\|. \tag{6.21}$$

Since

$$\|P_{\mathcal{M}^\perp} - P_{\mathcal{N}^\perp}\| = \|(I - P_{\mathcal{M}}) - (I - P_{\mathcal{N}})\| = \|P_{\mathcal{M}} - P_{\mathcal{N}}\|$$

hence

$$\theta(\mathcal{M}^\perp, \mathcal{N}^\perp) = \theta(\mathcal{M}, \mathcal{N}),$$

together with (6.21) and (6.17) imply the desired (6.18). \square

6.4. Several useful bounds

Let λ be an eigenvalue of $A_0 \in \mathbb{C}^{n \times n}$. Recall that the root subspace $\mathcal{R}(A_0, \lambda)$ is defined as a linear span of all Jordan chains of A_0 corresponding to λ , see, e.g., [5]. Alternatively, $\mathcal{R}(A_0, \lambda) = \text{Ker}(A_0 - \lambda I)^n$. Clearly, the dimension of $\mathcal{R}(A_0, \lambda)$ is equal to the total algebraic multiplicity of the eigenvalue λ of A_0 .

Further, let Γ be a simple (without self-intersections), closed, rectifiable contour with no eigenvalues of A_0 on it. Let $\{\lambda_1, \dots, \lambda_\gamma\}$ be a set of all eigenvalues of A_0 inside Γ . Denote

$$\mathcal{R}(A_0, \Gamma) = \mathcal{R}(A_0, \lambda_1) \dot{+} \dots \dot{+} \mathcal{R}(A_0, \lambda_\gamma).$$

With these notations the following bound holds.

Lemma 6.11 (Bound for the perturbation of root subspaces). *Let $A_0 \in \mathbb{C}^{n \times n}$, and let Γ be a simple, closed, rectifiable contour such that A_0 does not have eigenvalues on Γ . Then there are constants $K_{\text{root}}, \delta > 0$ (depending on A_0 and Γ only) such that any matrix A satisfying $\|A - A_0\| \leq \delta$ and does not have any eigenvalues on Γ , then*

$$\theta(\mathcal{R}(A_0, \Gamma), \mathcal{R}(A, \Gamma)) \leq K_{\text{root}} \|A - A_0\|. \tag{6.22}$$

In particular, the total multiplicity of all eigenvalues inside Γ is the same for A_0 and A .

The proof for the latter lemma can be found, e.g., in [6, p. 334]. We will also need the following result (cf., e.g., with [14]).

Lemma 6.12 (Bound for adjusting matrix S). *Let the decomposition*

$$\mathbb{C}^n = \mathcal{M}_1 \dot{+} \mathcal{M}_2 \dot{+} \dots \dot{+} \mathcal{M}_k \tag{6.23}$$

be given. For any decomposition

$$\mathbb{C}^n = \mathcal{N}_1 \dot{+} \mathcal{N}_2 \dot{+} \dots \dot{+} \mathcal{N}_k \tag{6.24}$$

with

$$\dim \mathcal{M}_j = \dim \mathcal{N}_j, \quad j = 1, 2, \dots, k, \tag{6.25}$$

there exists an invertible matrix $S \in \mathbb{C}^{n \times n}$ (that we suggest to call the adjusting matrix) satisfying

$$S\mathcal{M}_j = \mathcal{N}_j, \tag{6.26a}$$

$$\|I - S\| \leq \sum_{j=1}^k \theta(\mathcal{M}_j, \mathcal{N}_j). \tag{6.26b}$$

Proof. The proof is presented by considering two cases.

Case 1. $\theta(\mathcal{M}_j, \mathcal{N}_j) < 1$, for $j = 1, 2, \dots, k$. It is easy to verify that the matrix S defined as

$$S = I - \sum_{j=1}^k (P_{\mathcal{M}_j} - P_{\mathcal{N}_j}), \tag{6.27}$$

satisfy

$$S\mathcal{M}_j \subset \mathcal{N}_j. \tag{6.28}$$

Secondly, S is invertible. Indeed, for any nonzero $x \in \mathbb{C}^n$, let $x = x_1 + x_2 + \dots + x_k$ where $x_j \in \mathcal{M}_j$ we have

$$Sx = P_{\mathcal{N}_1}x_1 + \dots + P_{\mathcal{N}_k}x_k.$$

Since the $\theta(\mathcal{M}_j, \mathcal{N}_j) < 1$, hence \mathcal{M}_j and \mathcal{N}_j are not orthogonal. Therefore, if $x_j \neq 0$ then $P_{\mathcal{N}_j}x_j \neq 0$. This implies $\ker S = \{0\}$ so that S is invertible. In view of (6.25), the invertibility of S and (6.28) imply (6.26a). For S defined by (6.27) the relation (6.26b) is obvious:

$$\|I - S\| \leq \sum_{j=1}^k \|P_{\mathcal{N}_j} - P_{\mathcal{M}_j}\| = \sum_{j=1}^k \theta(\mathcal{M}_j, \mathcal{N}_j).$$

Case 2. $\theta(\mathcal{M}_j, \mathcal{N}_j) = 1$, for some j .

In view of (6.23), (6.24) and (6.25) we can always choose an invertible matrix T such that $T\mathcal{M}_j = \mathcal{N}_j$. Setting $S = \eta(\|T\|)^{-1}T$, where $\eta = \min_j \theta(\mathcal{M}_j, \mathcal{N}_j)$, we have

$$\|I - S\| \leq 1 + \eta \leq \sum_{j=1}^k \theta(\mathcal{M}_j, \mathcal{N}_j)$$

as desired. \square

To obtain necessary bounds on the perturbation of the eigenvalues of matrices we will need the following auxiliary result.

Lemma 6.13. *Let $A_0 \in \mathbb{C}^{n \times n}$, $\lambda \in \mathbb{C}$, and $a_1, a_2 \in \mathbb{C}^n$ satisfy $(A_0 - \lambda I)a_2 = a_1$. There is a constant $\delta > 0$ (depending on A_0 , a_1 , and a_2 only) such that for any $A \in \mathbb{C}^{n \times n}$, $\mu \in \mathbb{C}$, and $b_1, b_2 \in \mathbb{C}^n$ satisfying $(A - \mu I)b_2 = b_1$ and*

$$\|A - A_0\| \leq \delta, \quad |\mu - \lambda| \leq K_{\text{eig}}\|A - A_0\|, \quad \|b_2 - a_2\| \leq K_{\text{vec}}\|A - A_0\| \tag{6.29}$$

with some $K_{\text{eig}}, K_{\text{vec}} > 0$ we have

$$\|b_1 - a_1\| \leq K_{\text{next}}\|A - A_0\| \tag{6.30}$$

with

$$K_{\text{next}} = \|A_0\| + \delta + \|a_2\|(K_{\text{eig}} + 1) + (K_{\text{eig}}\delta + |\lambda|)K_{\text{vec}} \tag{6.31}$$

Proof. It follows from the first two inequalities in (6.29) that

$$\|A\| \leq \|A_0\| + \delta, \quad |\mu| \leq K_{\text{eig}}\delta + |\lambda|.$$

Using this and all the bounds in (6.29) we have

$$\begin{aligned} \|b_1 - a_1\| &= \|(A - \mu I)b_2 - (A_0 - \lambda I)a_2\| \\ &= \|Ab_2 - Aa_2 + Aa_2 - A_0a_2 + \lambda a_2 - \mu a_2 + \mu a_2 - \mu b_2\| \\ &\leq \|A\| \cdot \|b_2 - a_2\| + \|a_2\| \cdot \|A - A_0\| + \|a_2\| \cdot |\mu - \lambda| + |\mu| \cdot \|b_2 - a_2\| \\ &\leq (\|A_0\| + \delta + \|a_2\|(K_{\text{eig}} + 1) + (K_{\text{eig}}\delta + |\lambda|)K_{\text{vec}})\|A - A_0\|, \end{aligned}$$

and (6.30) follows. \square

Finally, here is the second key lemma of this section to be used in the proof of Lemma 5.2 in Section 7.

Lemma 6.14 (Perturbations of the eigenvalues and of Jordan chains). *Let Γ be a simple, closed, rectifiable contour such that A_0 does not have eigenvalues on Γ . Let $\sigma_1(A_0) = \{\lambda_1, \dots, \lambda_\gamma\}$ be the set of all eigenvalues of $A_0 \in \mathbb{C}^{n \times n}$ inside Γ . There are constants $K, \delta > 0$ (depending on A_0 only) such that the following statements holds. For any $A \in \mathbb{C}^{n \times n}$ satisfying*

$$\|A - A_0\| \leq \delta, \tag{6.32}$$

we have the following.

- (i) *If A has exactly γ eigenvalues inside Γ , there is a certain ordering $\{\mu_1, \dots, \mu_\gamma\}$ of them such that*

$$|\lambda_i - \mu_i| \leq K\|A - A_0\|, \quad i = 1, 2, \dots, \gamma. \tag{6.33}$$

(ii) Let $\{f_k\}_{k=0}^{m-1}$ be the longest Jordan chain corresponding to the eigenvalue $\lambda_i \in \sigma_1(A_0)$. If the maximal length of the Jordan chain corresponding to its eigenvalue μ_k described in (6.33) is also m , then there exists a Jordan chain $\{g_k\}_{k=0}^{m-1}$ of A corresponding to μ_i such that

$$\|f_k - g_k\| \leq K \|A - A_0\|, \quad k = 0, \dots, m - 1. \tag{6.34}$$

Proof. Here is the proof of the two parts of the lemma.

(i) For $i = 1, 2, \dots, \gamma$ let Γ_i denote a small circle that contains only one eigenvalue λ_i of A_0 . In accordance with Lemma 6.11 there is $\delta > 0$ such that any matrix A satisfying $\|A - A_0\| \leq \delta$ will have at least one eigenvalue inside Γ_i (indeed, the total multiplicity of the eigenvalues inside each Γ_i is preserved). Since A has exactly γ eigenvalues hence it must have exactly one eigenvalue, say, μ_i inside each Γ_i . Using Lemma 6.11 again we see that there are constants $K_i > 0$ (depending only on A_0) such that

$$\theta(\mathcal{R}(A_0, \lambda_i), \mathcal{R}(A, \mu_i)) \leq K_i \|A - A_0\|, \quad i = 1, \dots, \gamma. \tag{6.35}$$

Denote

$$\mathcal{M}_i = \mathcal{R}(A_0, \lambda_i), \quad \mathcal{N}_i = \mathcal{R}(A, \mu_i), \quad i = 1, \dots, \gamma,$$

and

$$\mathcal{M}_{\gamma+1} = \mathcal{N}_{\gamma+1} = (\mathcal{M}_1 \dot{+} \dots \dot{+} \mathcal{M}_\gamma)^\perp.$$

By Lemma 6.12 there is an S satisfying

$$S\mathcal{R}(A_0, \lambda_i) = \mathcal{R}(A, \mu_i), \quad i = 1, \dots, \gamma, \tag{6.36}$$

and

$$\|I - S\| \leq \sum_{i=1}^{\gamma} \theta(\mathcal{R}(A_0, \lambda_i), \mathcal{R}(A, \mu_i)). \tag{6.37}$$

Combining the latter two bounds (6.35) and (6.37) one obtains

$$\|I - S\| \leq K_0 \|A - A_0\| \tag{6.38}$$

with $K_0 = K_1 + \dots + K_\alpha$.

Now, let matrix R be such that

$$R^{-1}A_0R = \left[\begin{array}{ccc|c} A_1^{(0)} & 0 & \dots & * \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & A_\gamma^{(0)} & * \\ \hline 0 & \dots & 0 & A_{\gamma+1}^{(0)} \end{array} \right],$$

where $A_i^{(0)}$ has the only eigenvalue $\lambda_i \in \sigma_1(A_0)$, and all the eigenvalues of $A_{\gamma+1}^{(0)}$ are outside of Γ . Observe that R depends on A_0 only, and it can be fixed in advance. The property (6.36) yields that R diagonalizes $A_1 = S^{-1}AS$ as well, and moreover,

$$R^{-1}A_1R = \left[\begin{array}{ccc|c} A_1^{(1)} & 0 & \dots & * \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & A_\gamma^{(1)} & * \\ \hline 0 & \dots & 0 & A_{\gamma+1}^{(1)} \end{array} \right]$$

with each $A_i^{(1)}$ having only one eigenvalue μ_i , and with $A_{\gamma+1}^{(1)}$ having only the eigenvalues outside of Γ .

Further, denoting the size of $A_i^{(0)}$ by r_i we have

$$\begin{aligned} |\lambda_i - \mu_i| &\leq \frac{1}{r_i} |\text{trace}(A_i^{(0)}) - \text{trace}(A_i^{(1)})| \\ &= \frac{1}{r_i} \left| \sum_{k=1}^{r_i} e_k^T (A_i^{(0)} - A_i^{(1)}) e_k \right| \leq \|A_i^{(0)} - A_i^{(1)}\| \\ &\leq \|R^{-1} A_0 R - R^{-1} A_1 R\| \leq \kappa(R) \|A_0 - A_1\| \\ &= \kappa(R) \|A_0 - S^{-1} A S\| \leq K \|A_0 - A\|. \end{aligned}$$

In the above chain of inequalities the latter one is deduced from (6.38) using the arguments identical to those of the proof of Lemma 4.8. This concludes the proof of the part (i).

(ii) Let us define

$$g_{m-1} = P_{\mathcal{R}(A)} f_{m-1}, \quad g_{k-1} = (A - \mu_i I) g_k, \quad k = 1, \dots, m - 1.$$

It follows from (6.35) that

$$\begin{aligned} \|f_{m-1} - g_{m-1}\| &= \|f_{m-1} - P_{\mathcal{R}(A)} f_{m-1}\| \\ &\leq \|f_{m-1}\| \cdot \theta(\mathcal{R}(A_0, \lambda_i), \mathcal{R}(A, \mu_i)) \leq K_{\text{vec}} \|A - A_0\|, \end{aligned}$$

where $K_{\text{vec}} = \|f_{m-1}\| K_j$. Lemma 6.13 implies that

$$\|f_{m-2} - g_{m-2}\| \leq K_{\text{next}} \|A - A_0\|,$$

with K_{next} given by (6.31). It is easy to see from (6.31) that since K_{vec} depends on A_0 and the choice of the (fixed) chain $\{f_k\}$ hence the constant K_{next} , while possibly bigger, has the same property. Applying the same arguments recursively to f_{m-2}, g_{m-2} , then to f_{m-3}, g_{m-3} , and so on, one obtains, after m steps, the desired bound (6.34), in which the constant K is the maximum of the constants obtained in each of these steps.

To complete the proof of (ii) we need to show that $\{g_k\}_{k=0}^{m-1}$ is indeed a Jordan chain. To this end we need to show two things.

First, the vectors $\{g_k\}_{k=0}^{m-1}$ have to be linearly independent. Since $\delta > 0$ is at our disposal we may assume it to be small enough so that the bound (6.32), linear independence of $\{f_k\}$ and (6.34) guarantee linear independence of $\{f_k\}$.

Secondly, we need to show that $(A - \mu_i I) g_0 = 0$. This follows from our assumption that the longest Jordan chain of A corresponding to μ_i has length m .

7. Proof of Lemma 5.2

In this section we prove Lemma 5.2 which completes the proof of Theorem 5.1 and thus of the main result, Theorem 1.8. As before, we start with a clarifying example.

7.1. Multiple Block Model example

For some $\delta > 0$, consider the matrices A_0 and H_0 defined by

$$A_0 = \left[\begin{array}{ccc|cc} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \quad \text{and} \quad H_0 = \left[\begin{array}{ccc|cc} 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{array} \right],$$

and small perturbations of the above matrices A and H as,

$$A = \left[\begin{array}{ccc|cc} 0 & 1 & 2\delta & 0 & \delta \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \quad \text{and} \quad H = \left[\begin{array}{ccc|cc} 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & \delta \\ 1 & 0 & 2\delta & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 1 \\ 0 & \delta & 0 & 1 & 0 \end{array} \right].$$

We seek a matrix S such that

$$(A, H) \xrightarrow{S} (A_1, H_1), \quad \|I - S\| \leq K(\|A - A_0\| + \|H - H_0\|),$$

where A_1 and H_1 have the forms

$$A_1 = \left[\begin{array}{ccc|cc} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & * & * \\ 0 & 0 & 0 & * & * \end{array} \right] \quad \text{and} \quad H_1 = \left[\begin{array}{ccc|cc} 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & * & * \\ 0 & 0 & 0 & * & * \end{array} \right];$$

that is, the matrix S decouples the first Jordan chain from the remaining ones. Such a process in general enables us to proceed inductively on the later portions. As in Section 4.1 the matrix S is found in two steps, the first is such that

$$(A, H) \xrightarrow{S_1} (A_1, H'_1) \quad \text{and} \quad \|I - S_1\| \leq K(\|A - A_0\| + \|H - H_0\|),$$

where H'_1 has the form

$$H'_1 = \left[\begin{array}{ccc|cc} 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & * & 0 & 0 \\ 1 & * & * & 0 & 0 \\ \hline 0 & 0 & 0 & * & * \\ 0 & 0 & 0 & * & * \end{array} \right].$$

Then, in the second step, we produce a matrix S_2 such that

$$(A_1, H'_1) \xrightarrow{S_2} (A_1, H_1) \quad \text{and} \quad \|I - S_2\| \leq K(\|A - A_0\| + \|H - H_0\|).$$

Finally, we show that $S = S_1 S_2$ satisfies

$$(A, H) \xrightarrow{S} (A_1, H_1) \quad \text{and} \quad \|I - S\| \leq K(\|A - A_0\| + \|H - H_0\|)$$

as desired.

7.1.1. First step. Mapping $A \rightarrow A_1$. Constructing S_1

As a first attempt, we will try to proceed as in the example in Section 4.1 and choose a similarity matrix S_1 that maps the first Jordan chain of A_0 ,

$$0 \leftarrow e_1 \leftarrow e_2 \leftarrow e_3,$$

to that of A ,

$$0 \leftarrow e_1 \leftarrow e_2 + (2\delta)e_1 \leftarrow e_3.$$

This attempt will fail, but it will indicate a difficulty and a way to resolve it. Denote these vectors by g_k ,

$$g_3 = e_3, \quad g_2 = e_2 + (2\delta)e_1, \quad g_1 = e_1.$$

As before we choose an S_1 such that $S_1 : e_k \rightarrow g_k$ for $k = 1, 2, 3$, but this leaves a choice of where to map the vectors in root subspaces corresponding to later Jordan chains, in this case e_4 and e_5 . As a first attempt, let us choose S_1 to leave these unchanged. So we initially choose to map them as follows:

$$\begin{array}{l} S_1 : e_1 \rightarrow g_1 \\ S_1 : e_2 \rightarrow g_2 \\ S_1 : e_3 \rightarrow g_3 \\ S_1 : e_4 \rightarrow e_4 \\ S_1 : e_5 \rightarrow e_5 \end{array} \Leftrightarrow S_1 = \left[\begin{array}{ccc|cc} 1 & 2\delta & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ \hline 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right].$$

This choice fails to satisfy our requirements, as

$$S_1^{-1}AS_1 = \left[\begin{array}{ccc|cc} 0 & 1 & 0 & 0 & \delta \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \quad \text{and} \quad S_1^*HS_1 = \left[\begin{array}{ccc|cc} 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2\delta & 0 & \delta \\ 1 & 2\delta & 2\delta & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 1 \\ 0 & \delta & 0 & 1 & 0 \end{array} \right]$$

have nonzero off diagonal blocks. The resolution is to choose S_1 to map not only the vectors of the first chain of A_0 to that of A , but also map the other chains appropriately. Define

$$g_4 = [0 \quad 0 \quad 0 \quad 1 \quad 0]^T, \quad g_5 = [0 \quad -\delta \quad 0 \quad \delta^2 \quad 1]^T.$$

It is straightforward to check that $[x, y]_H = 0$ for $x \in \text{span}\{g_1, g_2, g_3\}$ and $y \in \text{span}\{g_4, g_5\}$; that is, the first root subspace $\text{span}\{g_1, g_2, g_3\}$ is H -orthogonal to all other root subspaces, in this case $\text{span}\{g_4, g_5\}$. Choosing the matrix S_1 such that

$$\begin{array}{l} S_1 : e_1 \rightarrow g_1 \\ S_1 : e_2 \rightarrow g_2 \\ S_1 : e_3 \rightarrow g_3 \\ S_1 : e_4 \rightarrow g_4 \\ S_1 : e_5 \rightarrow g_5 \end{array} \Leftrightarrow S_1 = \left[\begin{array}{ccc|cc} 1 & 2\delta & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & -\delta \\ 0 & 0 & 1 & 0 & 0 \\ \hline 0 & 0 & 0 & 1 & \delta^2 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right]$$

results in

$$S_1^{-1}AS_1 = \left[\begin{array}{ccc|cc} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \quad \text{and} \quad S_1^*HS_1 = \left[\begin{array}{ccc|cc} 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2\delta & 0 & 0 \\ 1 & 2\delta & 2\delta & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & \delta^2 \end{array} \right]$$

and so

$$(A, H) \xrightarrow{S_1} (A_1, H_1),$$

and since all perturbations are of the order δ , it is easy to see the required bound is satisfied. Hence S_1 is a correct choice in that it decouples and allows us to proceed inductively.

7.1.2. *Second step. Zeroing sub-antidiagonal entries of H_1 . Constructing S_2*

In the second step we produce a matrix S_2 that eliminates the 2δ sub-antidiagonal elements of the upper left submatrix of H_1 to produce the desired structure. Define

$$S_2 = \left[\begin{array}{ccc|cc} 1 & -\delta & \frac{3}{2}\delta^2 - \delta & 0 & 0 \\ 0 & 1 & -\delta & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ \hline 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right],$$

so that

$$S_2^{-1} S_1^{-1} A S_1 S_2 = \left[\begin{array}{ccc|cc} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right], \quad S_2^* S_1^* H S_1 S_2 = \left[\begin{array}{ccc|cc} 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & \delta^2 \end{array} \right].$$

7.1.3. *Third step. Combining affiliation matrices S_1 and S_2*

Define the matrix $S = S_1 S_2$, and so

$$S = \left[\begin{array}{ccc|cc} 1 & -\delta & \frac{3}{2}\delta^2 - \delta & 0 & 0 \\ 0 & 1 & -\delta & 0 & -\delta \\ 0 & 0 & 1 & 0 & 0 \\ \hline 0 & 0 & 0 & 1 & \delta^2 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right]$$

can easily be seen to satisfy the bound as all differences from identity are on the order of δ . The previous steps have shown that

$$(A, H) \xrightarrow{S_1} (A_1, H'_1) \xrightarrow{S_2} (A_1, H_1)$$

and so

$$(A, H) \xrightarrow{S} (A_1, H_1)$$

of the desired form. With the first block decoupled, we can proceed inductively.

7.2. *Proof of Lemma 5.2*

Part (i) follows from Lemma 6.14. Let us prove part (ii). Following the structure of the above example we prove Lemma 5.2 in three steps.

7.2.1. *First Step. Mapping $A \rightarrow A_1$. Constructing S_1*

In this section we prove the following proposition.

Proposition 7.1 (First decoupling step). *Let $A_0 \in \mathbb{C}^{n \times n}$ be a fixed H_0 -selfadjoint matrix, both A_0 and H_0 in the canonical form described in Theorem 2.1, and assume that A_0 has a real eigenvalue λ_1 . Denote*

$$A_0 = \begin{bmatrix} J(\lambda_1) & 0 \\ 0 & \tilde{A}_0 \end{bmatrix} \quad \text{and} \quad H_0 = \begin{bmatrix} \epsilon_1 P_1 & 0 \\ 0 & \tilde{H}_0 \end{bmatrix}; \tag{7.1}$$

i.e. consider the partition that singles out the largest, say, $m \times m$, Jordan block $J(\lambda_1)$ of A_0 . Then there are positive constants K and δ (depending on A_0 and H_0 only) such that for any H -selfadjoint matrix A with the same Jordan structure as A_0 satisfying

$$\|A - A_0\| + \|H - H_0\| < \delta,$$

the following statements hold.

(i) A has an eigenvalue μ_1 satisfying

$$|\lambda_1 - \mu_1| \leq K \|A - A_0\|. \tag{7.2}$$

(ii) There exists an affiliation matrix S_1 , i.e., $(A, H) \xrightarrow{S_1} (A_1, H_1)$ such that matrices A_1 and H_1 have the form

$$A_1 = \begin{bmatrix} J(\mu_1) & 0 \\ 0 & \tilde{A}_1 \end{bmatrix} \quad \text{and} \quad H_1 = \begin{bmatrix} \tilde{R}_1 & 0 \\ 0 & \tilde{H}_1 \end{bmatrix} \tag{7.3}$$

with some $\mu_1, \tilde{A}_1, \tilde{R}_1, \tilde{H}_1$, and S_1 satisfies the bound

$$\|I - S_1\| \leq K (\|A - A_0\| + \|H - H_0\|). \tag{7.4}$$

Proof. (i) Since $J(\lambda_1)$ is the largest Jordan block of A_0 corresponding to λ_1 , part (i) follows from Lemma 6.14.

(ii) The proof of (ii) will be based on the following procedure.

Procedure to construct the matrix S_1 of the Proposition 7.1

- Defining f_k 's, \mathcal{F}_1 and \mathcal{F}_2 . Since A_0 of Proposition 7.1 has canonical Jordan form, we can set $f_k = e_k$, the standard basis vectors. Further, set

$$\mathcal{F}_1 = \text{span}\{f_1, \dots, f_m\}, \quad \mathcal{F}_2 = \text{span}\{f_{m+1}, \dots, f_n\},$$

where m is the size of the largest Jordan block $J(\lambda)$ of A_0 in (7.1).

- Defining g_k 's, \mathcal{G}_1 and \mathcal{G}_2 . Let

$$\mathcal{G}_1 = \text{span}\{g_1, \dots, g_m\},$$

where $\{g_k\}_{k=1}^m$ are those described in Lemma 6.14 and satisfying (6.34). Now, in order to define the rest of the vectors $\{g_k\}_{k=m+1}^n$ we need to make several observations.

– First, $\mathcal{F}_1 \perp_{H_0} \mathcal{F}_2$. This follows easily from the block-diagonal structure of H_0 in (7.1) and the definition of indefinite inner product.

– Second, by Lemma 6.8 and (6.34) we have $\theta(\mathcal{F}_1, \mathcal{G}_1) \leq K_0 \|A - A_0\|$, where the constant $K_1 > 0$ depends on A_0 only and can be fixed in advance. Hence, if we define

$$\mathcal{G}_2 = \mathcal{G}_1^{[\perp]H},$$

then by Lemma 6.10 we will have

$$\theta(\mathcal{F}_2, \mathcal{G}_2) \leq K_1 (\|A - A_0\| + \|H - H_0\|).$$

Therefore, if for $k = m + 1, \dots, n$ we define $g_k = P_{\mathcal{G}_2} f_k$ then

$$\|f_i - g_i\| \leq K (\|A - A_0\| + \|H - H_0\|). \tag{7.5}$$

Again, the analysis shows that the constant $K > 0$ depends on A_0 and H_0 only and can be fixed in advance.

- *Defining S_1 .* Let us define $S_1 : f_k \rightarrow g_k$ for $k = 1, \dots, n$.

We are now ready to complete the proof of (ii). It is easy to see that the bounds (6.34) and (7.5) imply the desired bound (7.4), the proof of it follows along the lines of the proof of Proposition 4.2, literally. Hence it remains only to show that S_1 yields via $(A, H) \xrightarrow{S_1} (A_1, H_1)$ matrices A_1 and H_1 having the structure shown in (7.3).

First, recall that $S_1 : \mathcal{F}_k \rightarrow \mathcal{G}_k$. This fact and $H_1 = S_1^* H S$ allow us to deduce

$$\mathcal{F}_1[\perp]_{H_1} \mathcal{F}_2 \tag{7.6}$$

from $\mathcal{G}_1[\perp]_H \mathcal{G}_2$. Relation (7.6) means that in the standard basis $\{f_k\}$ the matrix H_1 has the block diagonal form shown in (7.3).

Secondly, $S_1 : f_k \rightarrow g_k$ means that S_1 maps the Jordan chain $\{f_k\}_{k=1}^m$ of A_0 to a Jordan chain $\{g_k\}_{k=1}^m$ of $A_1 = S_1^{-1} A S_1$. Hence \mathcal{F}_1 is A_1 -invariant, and moreover, the first block column of A_1 must have the form shown in (7.3). Finally, it is well-known that an H_1 -orthogonal complement \mathcal{F}_2 of an A_1 -invariant subspace \mathcal{F}_1 must be also A_1 -invariant, see, e.g. [6]. Hence, the second block column of A_1 must have the form shown in (7.3), and the proposition is completely proved. \square

Remark 7.2 (*First step in the case when A_0 has only complex eigenvalues*). In this case one has to consider the partitioning

$$A_0 = \begin{bmatrix} J_k(\lambda) & 0 \\ 0 & J_k(\bar{\lambda}) \end{bmatrix}, \quad H_0 = \tilde{I}$$

and the proof follows the same lines as above (as it was in Section 4.3).

7.2.2. Second Step. Zeroing sub-antidiagonal entries of H_1 . Constructing S_2 .

In the first step above we have constructed S_1 and mapped $(A, H) \xrightarrow{S_1} (A_1, H_1)$ shown in (7.3). Here we proceed with the upper left blocks $(J(\mu), \tilde{R}_1)$ of (A_1, H_1) and construct \tilde{S}_2 such that $(J(\mu), \tilde{R}_1) \xrightarrow{\tilde{S}_2} (J(\mu), \epsilon P_1)$. Theorem 4.1 implies that such \tilde{S}_2 exists and it must satisfy

$$\|I - \tilde{S}_2\| \leq K \|\tilde{R}_1 - \epsilon P_1\|.$$

The latter relation yields

$$\|I - S_2\| \leq K \|H_1 - H_0\|, \tag{7.7}$$

where $S_2 = \begin{bmatrix} \tilde{S}_2 & 0 \\ 0 & I \end{bmatrix}$ satisfy $(A_1, H_1) \xrightarrow{S_2} (A_2, H_2)$.

7.2.3. Third Step. Combining the affiliation matrices S_1 and S_2

In the two previous steps we have constructed matrices S_1 and S_2 such that

$$(A, H) \xrightarrow{S_1} (A_1, H_1) \xrightarrow{S_2} (A_2, H_2).$$

Hence $S = S_1 S_2$ satisfies $(A, H) \xrightarrow{S} (A_2, H_2)$ and the desired bound (5.5) now follows from (7.4), (7.7), and Theorem 4.9.

This completes the proof of Lemma 5.2.

8. Proof of Theorem 2.6. Global stability of sign characteristic. Weak affiliation matrix vs strong affiliation matrix

The last result to prove in the flow chart of Section 2.5 is Theorem 2.6. We will prove it in Section 8.1. As a corollary, we will then deduce in Section 8.2 a classical result of [4] on the stability of sign characteristic. Both results will be needed in Section 8.3 to define the concept of a strong affiliation matrix.

8.1. Proof of Theorem 2.6

First, by the definition of a canonical Jordan basis, the (fixed) vectors

$$\left\{ \left\{ f_r^{(k,s)} \right\}_{r=0}^{m_k(A_0, \lambda_s) - 1} \right\}_{s=1, k=1}^{s=\beta, k=\dim \text{Ker}(A_0 - \lambda_s I)}$$

are the columns of a certain (fixed) affiliation matrix F (i.e., satisfying

$$(A_0, H_0) \xrightarrow{F} (J_0, P_0), \tag{8.1}$$

where (J_0, P_0) is canonical). Indeed, it follows from

$$F^{-1} A_0 F = J_0.$$

Secondly, A and A_0 have the same Jordan structure and hence by Theorem 1.8 there exists S satisfying (1.7) and such that

$$(A, H) \xrightarrow{S} (A_1, H_0), \tag{8.2}$$

where the matrix

$$A_1 = S^{-1} A S$$

has the same Jordan bases as A_0 . So,

$$(A_1, H_0) \xrightarrow{F} (J_1, P_0), \tag{8.3}$$

where J_1 is Jordan. In particular, we have that

$$F^{-1} A_1 F = J_1.$$

It follows that

$$F^{-1} S^{-1} A S F = J_1.$$

Denoting by

$$\left\{ \left\{ g_r^{(k,s)} \right\}_{r=0}^{m_k(A, \mu_s) - 1} \right\}_{s=1, k=1}^{s=\beta, k=\dim \text{Ker}(A - \mu_s I)}$$

the columns of the matrix

$$G := S F, \tag{8.4}$$

we see that both bases $\{f_r^{(k,s)}\}$ and $\{g_r^{(k,s)}\}$ are canonical, and the desired (2.8) is, in fact, a reformulation of (1.7). \square

To introduce in Section 8.5 the concept of a (strong) affiliation matrix (as opposed to a weak affiliation matrix of Definition 1.7) we will need the following observation.

Remark 8.1. Let S be a (weak) affiliation matrix constructed in the process of the proof of Theorem 1.8. Then we have the following two observations.

(i) Formula (8.4) implies the following useful property of S :

$$g_r^{(k,s)} = S f_r^{(k,s)} \quad \text{for all } k, s, r \text{ within their ranges.} \tag{8.5}$$

(ii) Using the above notations, let the chains $\{f_r^{(k,1)}\}$ correspond to the eigenvalue λ_1 of A_0 , and let the chains $\{g_r^{(k,1)}\}$ correspond to the eigenvalue μ_1 of A . The flow chart for the proof of Theorem 1.8 in Section 2.5 indicates that Lemma 5.2 was crucial in constructing S . Therefore, by inspecting (5.3) (and also (7.2)) we see that the eigenvalue μ_1 of A had been chosen to be a small perturbation

$$|\lambda_1 - \mu_1| \leq K \|A - A_0\|,$$

of the eigenvalue λ_1 of A_0 .

The concept of a strong affiliation matrix will be introduced in Section 8.3 using two observations. One of them is the above Remark 8.1, and the second one is one classical result [4] on the stability of sign characteristic recalled next.

8.2. Stability of sign characteristic as a consequence of the stability of canonical bases

We begin this section with the following simple example.

Example 8.2 (Computing sign characteristic from a canonical Jordan basis). Let us return to the example 2.5 and examine one way of computing the sign characteristic of (J, P) from its canonical basis. In this case $\{e_1, e_2, e_3\}$ and $\{e_3, e_4\}$ are the two Jordan chains of J , and it is easy to see that

$$\epsilon_1 = 1 = [e_1, e_3]_P, \quad \epsilon_2 = -1 = [e_4, e_5]_P.$$

Theorem 2.1 and the argument similar to the one in the above example lead to the following obvious statement.

Lemma 8.3 (Computing the sign characteristic). *Let $A \in \mathbb{C}^{n \times n}$ be a fixed H -selfadjoint matrix. Let*

$$\left\{ \left\{ f_r^{(k,s)} \right\}_{r=0}^{m_k(A, \lambda_s) - 1} \right\}_{s=1, k=1}^{s=\beta, k=\dim \text{Ker}(A - \lambda_s I)} \tag{8.6}$$

and let $\{\lambda_1, \dots, \lambda_\alpha\}$ be all real eigenvalues of A be a fixed canonical Jordan basis of (A, H) . Then the sign characteristic (2.5) satisfies

$$\epsilon_{k,s} = [f_0^{(k,s)}, f_{m_k(A, \lambda_s) - 1}^{(k,s)}]_H, \quad (k = 1, \dots, \dim \text{Ker}(A - \lambda_s I), s = 1, \dots, \alpha).$$

It is easy to see that Theorem 2.6, part (ii) of Remark 8.1 and Lemma 8.3 imply the following well-known result [4,6,16].

Theorem 8.4 (Global stability of the sign characteristic). *Let $A_0 \in \mathbb{C}^{n \times n}$ be a fixed H_0 -selfadjoint matrix, and let $\{\lambda_1, \dots, \lambda_\alpha\}$ denote all distinct real eigenvalues of A_0 . Let $\gamma > 0$ be such that every real eigenvalue λ_k of A_0 is the only eigenvalue in the interval $(\lambda_k - \gamma, \lambda_k + \gamma)$. There*

exists a constant $\delta > 0$ (depending on A_0 , H_0 and γ only) such that the following assertion holds. For any H -selfadjoint matrix A such that A has the same Jordan structure as A_0 and

$$\|A - A_0\| + \|H - H_0\| < \delta,$$

matrix A has a unique eigenvalue, say, μ_k in the interval $(\lambda_k - \gamma, \lambda_k + \gamma)$ and the sign characteristics of λ_k and μ_k coincide (up to a rearrangement of the signs corresponding to the same block sizes).

The above result is *global*, i.e., it assumes that the Jordan structure of A_0 is preserved for *all* eigenvalues. In Corollary 9.6 of Section 9 we will also obtain a *local* version of this stability result.

8.3. Affiliation matrix revisited. Mapping canonical bases

Definition 1.7 introduced *weakly* affiliated matrices S . Moreover, up until this point the term “affiliated” was understood in the *weak* sense. However, it had been just observed in Section 8.2 that the affiliation matrix S had been constructed in such a way that it has several additional properties. We therefore define next an affiliation matrix as a weak affiliation matrix obeying those additional restrictions.

Definition 8.5 ((*Strong*) *affiliation matrix*). Let A_0 be H_0 -selfadjoint and A be H -selfadjoint. A matrix S is called a *strong affiliation matrix* (or just an *affiliation matrix*) of the quadruple (A_0, H_0, A, H) if there exist two canonical Jordan bases $\{f_r^{(k,s)}\}$ and $\{g_r^{(k,s)}\}$ of A_0 and A , respectively, such that

- (8.5) holds.
- S in (8.5) maps Jordan chains $\{f_r^{(k,s)}\}$ corresponding to real eigenvalues of A_0 to Jordan chains $\{g_r^{(k,s)}\}$ corresponding to real eigenvalues of A , and the same property holds for nonreal eigenvalues as well.
- The mapping $S : \{f_r^{(k,s)}\} \rightarrow \{g_r^{(k,s)}\}$ preserves the sign characteristic for each Jordan chain.

In this case pairs (A_0, H_0) and (A, H) are called (strongly) affiliated.

Remark 8.6 (*The difference between an affiliation relation and a weak affiliation relation*). We start with three obvious observations.

- Both weakly affiliated and strongly affiliated are equivalence relations.
- The equivalence class of all matrices that are strongly affiliated to a given pair (A, H) is a subset of all matrices that are weakly affiliated to (A, H) .
- Finally, each pair (A, H) is affiliated to its canonical form (J, P) described in Theorem 2.1.

Hence it is of interest to describe both weak and strong affiliation relations in term of canonical forms. The comparison of Definitions 1.7 and 8.5 yields the following facts.

- (i) Two pairs (A, H) and (B, G) (where A and B are H -selfadjoint and G -selfadjoint, respectively) are weakly affiliated if the matrices J_A and J_B (of their canonical forms (J_A, P_H) and (J_B, P_G)) have the same Jordan structure, i.e., there is a bijective function $f : \{\lambda_1, \dots, \lambda_\beta\}$

→ {μ₁, . . . , μ_β} such that J_B is obtained from J_A by replacing λ_k's by μ_k's. Here {λ₁, . . . , λ_β} and {μ₁, . . . , μ_β} are the sets of all eigenvalues of A and B, respectively.

There is no restriction on the structure of sip matrices P_H and P_G.

- (ii) Two pairs (A, H) and (B, G) are strongly affiliated if, in addition to the description of part (i), we also have P_H = P_G which means that not only A and B have same Jordan structure, but they also share the same sign characteristic.

Recall that in Theorem 1.8 we proved the existence of a weak affiliation matrix S. It is now clear that it is the stability of sign characteristic that allowed us to conclude that Theorem 1.8, in fact, claims the existence of a strong affiliation matrix S.

The “local” Definition 8.5 has, in certain circumstances, some advantages over the “global” Definition 1.7. For instance, in the next section it will be adapted to derive a variant of the main result for the case when the Jordan structure is preserved only for a selection of the eigenvalues.

9. Perturbations partially preserving Jordan structure

9.1. Balanced partitions

In this section, we present an extension of the case considered thus far to the case of perturbations that preserve the Jordan structure only for some selection of the eigenvalues. To be specific, let A₀ be H₀-selfadjoint, and let

$$\sigma(A_0) = \sigma_1(A_0) \cup \sigma_2(A_0) \quad (\text{with } \sigma_1(A_0) \cap \sigma_2(A_0) = \emptyset) \tag{9.1}$$

be a partition of the set σ(A₀) of all eigenvalues of A₀. Recall that in accordance with Remark 2.2, σ(A₀) is symmetric with respect to the real axis. In this section we consider only what we suggest to call *balanced partitions*, i.e., those for which σ₁(A₀) is symmetric about the real axis as well (in fact, σ₂(A₀) will be automatically symmetric as well in this case).

In the rest of the paper we extend the results of Sections 2–8 to the situation where Jordan structure is assumed to be preserved for the eigenvalues in σ₁(A₀) only.

9.2. Basic definitions. σ₁-Partial Jordan structure. σ₁-Partial affiliation matrix

In this section we provide a number of counterparts for the basic definitions and facts of Sections 1 and 2.

Definition 9.1 (A counterpart of Definition 1.2).

- (Same σ₁-partial Jordan structure). Let (9.1) be a balanced partition of the set of all eigenvalues of A₀. Matrices A₀ and A are said to have *the same σ₁-partial Jordan structure* if there is a balanced partition

$$\sigma(A) = \sigma_1(A) \cup \sigma_2(A) \tag{9.2}$$

and a bijective function f : σ₁(A₀) → σ₁(A) such that if μ = f(λ), then λ and μ have the same Jordan block sizes.

- (Same σ₁-partial Jordan bases). In this case, matrices A₀ and A are said to have *the same σ₁-partial Jordan bases* if the following statement is true. If μ = f(λ), then every Jordan chain

of A_0 corresponding to λ is also a Jordan chain of A corresponding to μ (and automatically vice versa).

The following statement is a counterpart of Remark 1.3

Remark 9.2 (Same σ_1 -partial Jordan bases). Let (9.1) and (9.2) be balanced partitions of A_0 and A , respectively. The matrices A_0 and A have the same Jordan σ_1 -partial bases if the following statement holds. If, for an invertible T , the matrix $T^{-1}A_0T$ has the form

$$T^{-1}A_0T = \begin{bmatrix} J_0 & * \\ 0 & M_0 \end{bmatrix}, \quad \text{with } \sigma(J_0) = \sigma_1(A_0), \sigma(M_0) = \sigma_2(A_0),$$

where J_0 is in a canonical Jordan form, then $T^{-1}AT$ has the form

$$T^{-1}AT = \begin{bmatrix} J_1 & * \\ 0 & M_1 \end{bmatrix}, \quad \text{with } \sigma(J_1) = \sigma_1(A), \sigma(M_1) = \sigma_2(A),$$

where J_1 is also in a canonical Jordan form.

The next definition is a counterpart of Definition 2.4.

Definition 9.3 (Canonical σ_1 -partial Jordan basis. σ_1 -Partial sign characteristic). Let A_0 be an H_0 -selfadjoint matrix. Let $\sigma(A_0) = \{\lambda_1, \dots, \lambda_\beta\}$ be the set of all eigenvalues of A_0 and let

$$\{\{f_r^{(k,s)}\}_{r=0}^{m_k(A_0,\lambda_s)-1}\}_{s=1,k=1}^{s=\beta,k=\dim \text{Ker}(A_0-\lambda_s I)} \tag{9.3}$$

be its canonical Jordan basis. Let (9.1) be a balanced partition with $\sigma_1(A_0) = \{\lambda_{j_1}, \dots, \lambda_{j_\gamma}\}$.

- The subset

$$\{\{f_r^{(k,j_s)}\}_{r=0}^{m_k(A_0,\lambda_{j_s})-1}\}_{s=1,k=1}^{s=\gamma,k=\dim \text{Ker}(A_0-\lambda_{j_s} I)} \tag{9.4}$$

of (9.3) is called a canonical σ_1 -partial Jordan basis of A_0 .

- A subset of signs in the sign characteristic of (A_0, H_0) corresponding to its canonical σ_1 -partial Jordan basis is called σ_1 -partial sign characteristic.

In words, a σ_1 -partial Jordan basis of A_0 is just a selection of its canonical Jordan chains corresponding to the eigenvalues belonging to the subset $\sigma_1(A_0)$, and the same is true for σ_1 -partial sign characteristic.

We are now ready to formulate a counterpart of Definition 8.5 and to define a “local” version of an affiliation matrix.

Definition 9.4 (σ_1 -Partial affiliation matrix). Let A_0 be H_0 -selfadjoint and A be H -selfadjoint. A matrix S is called a σ_1 -partial affiliation matrix of the quadruple (A_0, H_0, A, H) if there exist two σ_1 -partial canonical Jordan bases $\{f_r^{(k,s)}\}$ and $\{g_r^{(k,s)}\}$ of A_0 and A , respectfully, such that

- (8.5) holds.
- S in (8.5) maps Jordan chains $\{f_r^{(k,s)}\}$ corresponding to real eigenvalues in $\sigma_1(A_0)$ to Jordan chains $\{g_r^{(k,s)}\}$ corresponding to real eigenvalues in $\sigma_1(A)$, and the same property holds for nonreal eigenvalues in $\sigma_1(A_0)$ and $\sigma_1(A)$ as well.
- For each Jordan chain corresponding to real eigenvalues in $\sigma_1(A_0)$ the mapping $S : \{f_r^{(k,s)}\} \rightarrow \{g_r^{(k,s)}\}$ preserves the sign characteristic.

In this case pairs (A_0, H_0) and (A, H) are called σ_1 -partially affiliated.

9.3. Lipschitz stability of σ_1 -partial affiliation matrices

The main result of this section, an extension of Theorem 2.6 (and hence of Theorem 1.8), is stated next.

Theorem 9.5 (Lipschitz stability for σ_1 -partial perturbations). *Let $A_0 \in \mathbb{C}^{n \times n}$ be a fixed H_0 -selfadjoint matrix, and let (9.1) be a balanced partition for A_0 with $\sigma_1(A_0) = \{\lambda_1, \dots, \lambda_\gamma\}$. Let*

$$\left\{ \left\{ f_r^{(k,s)} \right\}_{r=0}^{m_k(A_0, \lambda_s) - 1} \right\}_{s=1, k=1}^{s=\gamma, k=\dim \text{Ker}(A - \lambda_s I)}$$

be a fixed σ_1 -partial canonical Jordan basis of A_0 . Finally, let Γ be a simple, closed rectifiable contour such that A_0 does not have eigenvalues on Γ and $\sigma_1(A_0)$ is the set of all eigenvalues of A_0 inside Γ .

There exist constants $K, \delta > 0$ (depending on A_0 and H_0 only) such that for any H -selfadjoint matrix A having the same σ_1 -partial Jordan structure as A_0 (where $\sigma_1(A)$ is the set of all eigenvalues of A inside Γ) and satisfying

$$\|A - A_0\| + \|H - H_0\| < \delta,$$

the following assertion holds.

- (i) (Lipschitz stability of σ_1 -partial affiliation matrices) *The pairs (A_0, H_0) and (A, H) are σ_1 -partially affiliated, and there exists a σ_1 -partial affiliation matrix S satisfying*

$$\|I - S\| \leq K (\|A - A_0\| + \|H - H_0\|). \tag{9.5}$$

- (ii) (Lipschitz stability of σ_1 -partial canonical Jordan bases) *There exists a σ_1 -partial canonical Jordan basis $\left\{ \left\{ g_r^{(k,s)} \right\}_{r=0}^{m_k(A, \mu_s) - 1} \right\}_{s=1, k=1}^{s=\gamma, k=\dim \text{Ker}(A - \mu_s I)}$ of A such that*

$$\|g_r^{(k,s)} - f_r^{(k,s)}\| \leq K (\|A - A_0\| + \|H - H_0\|) \tag{9.6}$$

for $k = 1, \dots, \gamma$ and all k, r within their ranges.

Before proving the above theorem in Section 9.4 we formulate the following obvious corollary.

Corollary 9.6 (Stability of σ_1 -partial sign characteristic). *Let $A_0 \in \mathbb{C}^{n \times n}$ be a fixed H_0 -selfadjoint matrix, and let (9.1) be a balanced partition for A_0 . There exist a constant $\delta > 0$ (depending on A_0 and H_0 only) such that for any H -selfadjoint matrix A having the same σ_1 -partial Jordan structure as A_0 and satisfying*

$$\|A - A_0\| + \|H - H_0\| < \delta,$$

the σ_1 -partial sign characteristics of A and A_0 coincide.

9.4. Proof of Theorem 9.5

Actually, the proof of Theorem 9.5 can be derived by just adapting all results of Sections 2–8 to the σ_1 -partial case. For example, here is a counterpart of the Theorem 3.1

Theorem 9.7 (Reduction to the canonical form). *Suppose the result of Theorem 9.5 is true for each pair (A_0, H_0) in the canonical form as defined in the Definition 2.3. Then the result of Theorem 9.5 is true for all pairs (B_0, G_0) , where B_0 is G -selfadjoint.*

Sketch of the proof. The proof follows the lines of the proof of Theorem 3.1 with only minor modifications. We therefore highlight only the major differences. Specifically, one has to modify the diagram (3.1) as follows.

$$\begin{array}{ccc}
 (A, H) & \xrightarrow{S} & (A_1, H_1) \\
 T \uparrow & & T \uparrow \\
 (B, G) & \xrightarrow{R=TSST^{-1}} & (B_1, G_1)
 \end{array} \tag{9.7}$$

Thus to replace H_0 and G_0 of (3.1) by H_1 and G_1 , respectfully. The point of this replacement is that if we partition the canonical pair (A_0, H_0) as

$$A_0 = \begin{bmatrix} A_1^{(0)} & 0 \\ 0 & A_2^{(0)} \end{bmatrix}, \quad H_0 = \begin{bmatrix} H_1^{(0)} & 0 \\ 0 & H_2^{(0)} \end{bmatrix}, \quad \text{with} \\
 \sigma(A_1^{(0)}) = \sigma_1(A_0), \sigma(A_2^{(0)}) = \sigma_2(A_0),$$

then the pair (A_1, H_1) should look like

$$A_1 = \begin{bmatrix} A_1^{(1)} & 0 \\ 0 & A_2^{(1)} \end{bmatrix}, \quad H_1 = \begin{bmatrix} H_1^{(1)} & 0 \\ 0 & H_2^{(1)} \end{bmatrix}, \quad \text{with} \\
 \sigma(A_1^{(1)}) = \sigma_1(A), \sigma(A_2^{(1)}) = \sigma_2(A),$$

where the pair $(A_1^{(1)}, H_1^{(1)})$ is canonical, i.e., $A_1^{(1)}$ is Jordan and $H_1^{(1)}$ is sip. Considering matrices (A_1, H_1) that have canonical structure only in their leading blocks is exactly what is needed for proving the desired result in the σ_1 -partial case.

The rest of the arguments follow the lines of the proof of Theorem 3.1.

Using the above result it is possible to complete the proof of Theorem 9.5 by adapting the rest of the proof of Theorem 5.1. Indeed, we proved Theorem 5.1 recursively, i.e., by decoupling Jordan blocks one by one. Adaptation of that proof to the σ_1 -partial case simply means stopping the decoupling process earlier, after processing all the eigenvalues in σ_1 .

However, instead of asking the reader to inspect the proof of Theorem 5.1 in Sections 4, 5, and 7 (see, e.g., the flow chart in Section 2.5) we prefer to give a short direct proof.

Proof of Theorem 9.5. The proof of (i) is the main part of the proof, the claim (ii) is just a corollary of (i).

(i) Let $\sigma_1(A_0) = \{\lambda_1, \dots, \lambda_\gamma\}$, and let the canonical pair (A_0, H_0) be partitioned as

$$A_0 = \begin{bmatrix} J^{(0)}(\lambda_1) & 0 & \cdots & 0 \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & J^{(0)}(\lambda_\gamma) & 0 \\ 0 & \cdots & 0 & A_{\gamma+1}^{(0)} \end{bmatrix}, \quad H_0 = \begin{bmatrix} P_1^{(0)} & 0 & \cdots & 0 \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & P_\gamma^{(0)} & 0 \\ 0 & \cdots & 0 & H_{\gamma+1}^{(0)} \end{bmatrix},$$

where each pair $(J^{(0)}(\lambda_i), P_i^{(0)})$ is in the canonical form, $\sigma(J^{(0)}(\lambda_i)) = \{\lambda_i\}$ and $\sigma(A_{\gamma+1}^{(0)}) \subset \sigma_2(A_0)$.

Further, let $\{\Gamma_i\}$ be a set of small non-intersecting circles such that λ_i is the only eigenvalue of A_0 inside each circle Γ_i . Since A has the same σ_1 -partial Jordan structure as A_0 there is $\delta > 0$ guaranteeing that A has only one eigenvalue, say, μ_i inside each Γ_i , i.e., $\sigma_1(A) = \{\mu_1, \dots, \mu_\gamma\}$. Denote

$$\mathcal{M}_i = \mathcal{R}(A_0, \lambda_i), \quad \mathcal{N}_i = \mathcal{R}(A, \mu_i), \quad i = 1, \dots, \gamma,$$

and

$$\mathcal{M}_{\gamma+1} = (\mathcal{M}_1 \dot{+} \dots \dot{+} \mathcal{M}_\gamma)^{[\perp]H_0}, \quad \mathcal{N}_{\gamma+1} = (\mathcal{N}_1 \dot{+} \dots \dot{+} \mathcal{N}_\gamma)^{[\perp]H},$$

Using Lemmas 6.10, 6.11, and 6.12 one can construct S_1 such that

$$S_1 \mathcal{M}_i = \mathcal{N}_i, \quad i = 1, \dots, \gamma + 1 \tag{9.8}$$

and

$$\|I - S_1\| \leq K_1(\|A - A_0\| + \|H - H_0\|) \tag{9.9}$$

for some K_1 depending on A_0 and H_0 only. Define (A_1, H_1) by

$$(A, H) \xrightarrow{S_1} (A_1, H_1).$$

Clearly, (9.8) and the fact that \mathcal{M}_i are H_0 -orthogonal and \mathcal{N}_i are H -orthogonal imply that the matrices in (A_1, H_1) have the same block form

$$A_1 = \begin{bmatrix} A^{(1)}(\mu_1) & 0 & \dots & 0 \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & A^{(1)}(\mu_\gamma) & 0 \\ 0 & \dots & 0 & A_{\gamma+1}^{(1)} \end{bmatrix}, \quad H_1 = \begin{bmatrix} H_1^{(1)} & 0 & \dots & 0 \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & H_\gamma^{(1)} & 0 \\ 0 & \dots & 0 & H_{\gamma+1}^{(1)} \end{bmatrix}$$

as (A, H) . Indeed, observe that at the moment the pairs $(A^{(1)}(\mu_i), H_i^{(1)})$ are not in the canonical form yet. Using Theorem 5.1, one finds matrices $S_i^{(2)}$ such that

$$(A^{(1)}(\mu_i), H_i^{(1)}) \xrightarrow{S_i^{(2)}} (J^{(1)}(\mu_i), P_i^{(1)}), \quad i = 1, 2, \dots, \gamma,$$

where the pairs $(J^{(1)}(\mu_i), P_i^{(1)})$ are canonical. Defining

$$S_2 = \begin{bmatrix} S_1^{(2)} & 0 & \dots & 0 \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & S_\gamma^{(2)} & 0 \\ 0 & \dots & 0 & I \end{bmatrix}$$

one sees that

$$(A_1, H_1) \xrightarrow{S_2} \left(\begin{bmatrix} J^{(1)}(\lambda_1) & 0 & \dots & 0 \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & J^{(1)}(\lambda_\gamma) & 0 \\ 0 & \dots & 0 & A_{\gamma+1}^{(1)} \end{bmatrix}, \begin{bmatrix} P_1^{(0)} & 0 & \dots & 0 \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & P_\gamma^{(0)} & 0 \\ 0 & \dots & 0 & H_{\gamma+1}^{(1)} \end{bmatrix} \right)$$

and by (5.1) we have

$$\|I - S_2\| \leq K_2(\|A_1 - A_0\| + \|H_1 - H_0\|). \quad (9.10)$$

Clearly, $S = S_1 S_1$ is the desired σ_1 -partial affiliation matrix. Finally, combining (9.9) and (9.10) and using Lemma 4.8 we obtain the bound (9.5).

(ii) The bound (9.5) and the relation $Sf_r^{(k,s)} = g_r^{(k,s)}$ yields the desired bound (9.6).

Acknowledgments

The authors would like to express their sincere appreciation and gratitude for the many helpful suggestions provided by Michael Neumann during the entire process of working on this manuscript. The quality of the paper has been improved as a result.

References

- [1] Rajendra Bhatia, *Matrix Analysis*, Springer, 1997.
- [2] H. Den Boer, G.Ph. Thijssse, Semi-stability of sums of partial multiplicities under additive perturbations, *Integral Equations Operator Theory* 3 (1980) 23–42.
- [3] I. Gohberg, M.A. Kaashoek, Unsolved problems in matrix and operator theory, *Integral Equations Operator Theory* 1, 278–283.
- [4] I. Gohberg, P. Lancaster, L. Rodman, *Matrices and Indefinite Scalar Products*, Birkhäuser, 1983.
- [5] I. Gohberg, P. Lancaster, L. Rodman, *Invariant Subspaces of Matrices with Applications*, Wiley-Interscience, 1986.
- [6] I. Gohberg, P. Lancaster, L. Rodman, *Indefinite Linear Algebra and Applications*, Birkhäuser, 2005.
- [7] I. Gohberg, P.L. Rodman, On distance between lattices of invariant subspaces of matrices, *Linear Algebra Appl.* 76 (1986) 85–120.
- [8] T. Kato, *Perturbation theory for linear operators*, Die Grundlehren der mathematischen Wissenschaften, Band 132 Springer-Verlag New York, Inc., New York, 1966 xix+592 pp. 47.00 (47.48).
- [9] M. Konstantinov, D. Gu, V. Mehrmann, P. Petkov, *Perturbation Theory for Matrix Equations Studies in Computational Mathematics*, Elsevier, North Holland, 2003.
- [10] A.I. Mal'cev, *Foundations of Linear Algebra*, in: W.H. Freeman (Ed.), San Francisco and London, 1963 (translation from Russian).
- [11] V. Matsaev, V. Olshevsky, Cyclic dimensions, kernel multiplicities and Gohberg-Kaashoek numbers, *Linear Algebra Appl.* 239 (1996) 161–174.
- [12] A. Markus, E. Parilis, The change of the Jordan structure of a matrix under small perturbations, *Mat. Issled.* 54 (1980) 98–109 (in Russian); A. Markus, E. Parilis, English translation: *Linear Algebra Appl.* 54 (1983) 139–152.
- [13] V. Olshevsky, A condition for the closeness of the sets of invariant subspaces of the close matrices in terms of their Jordan structures, *Siberian Math. J.* 30 (4) (1989) 102–110 (in Russian); V. Olshevsky, English translation: *Siberian Math. J.*, Vol. 30(4), Plenum Publishing Corp., 1989, pp. 580–586.
- [14] V. Olshevsky, Change of Jordan structure of G-selfadjoint operators and selfadjoint operator functions under small perturbations, *Izvestia Akad. Nauk USSR* 54 (5) 1990 1021–1048 (in Russian); V. Olshevsky, English translation: *AMS Math. USSR Izvestia* 37 (2) (1991) 371–396.
- [15] L. Rodman, S. Pierce, Congruences and norms of hermitian matrices, *Canad. Math. J.* XXXIX (1987) 1446–1458.
- [16] L. Rodman, Similarity vs unitary similarity and perturbation analysis of sign characteristics: Complex and real indefinite inner products, *Linear Algebra Appl.* 416 (2006) 945–1009.
- [17] G.W. Stewart, J.G. Sun, *Matrix Perturbation Theory*, Academic Press Inc., Boston, MA, 1990.
- [18] K. Weierstrass, Zur Theorie der quadratischen und bilinearen Formen, *Monatsber. Akad. Wiss., Berlin*, 1868, pp. 310–338.
- [19] K. Weierstrass, *Mathematische Werke, Zweiter Band. Abhandlungen II*. Berlin, Mayer & Muller, 1895, pp. 19–44.